

Rabbit PSMC2 Polyclonal Antibody | anti-PSMC2 antibody

PSMC2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA168225	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PSMC2; Polyclonal Antibody; PSMC2 Antibody; S7; MSS1; Nbla10058; anti-PSMC2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1mg/mL (varies by lot)
Sequence Length	433
Applicable Applications for anti-PSMC2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human PSMC2
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-74872 / sc-74874 / sc-133932 / sc-98691
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cell using PSMC2 antibody. Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human thyroid cancer using PSMC2 antibody at dilution of 1:200 (400x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human endometrial cancer using PSMC2 antibody at dilution of 1:200 (400x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PSMC2 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMC2 antibody

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit has been shown to interact with several of the basal transcription factors so, in addition to participation in proteasome functions, this subunit may participate in the regulation of transcription. This subunit may also compete with PSMC3 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

NCBI AND UNIPROT INFORMATION

NCBI GI #	649102561
NCBI Official Full Name	PSMC2, partial

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/psmc2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/168225>