

Rabbit ACVR2A Polyclonal Antibody | anti-ACVR2A antibody

ACVR2A Polyclonal Antibody

CATALOG # AAA168533	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

ACVR2A; Polyclonal Antibody; ACVR2A Polyclonal Antibody; ACVR2; ACTRII; anti-ACVR2A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2.9mg/mL (varies by lot)
Sequence Length	513
Applicable Applications for anti-ACVR2A antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Synthetic peptide of human ACVR2A
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.


IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human ovarian cancer using ACVR2A Polyclonal Antibody at dilution of 1:20



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human colon cancer using ACVR2A Polyclonal Antibody at dilution of 1:20



WB (Western Blot)

Western Blot analysis of Human fetal brain tissue using ACVR2A Polyclonal Antibody at dilution of 1:400

ADDITIONAL INFORMATION

Related Product Information for anti-ACVR2A antibody

This gene encodes activin A type II receptor. Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling, and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases.

Product Categories/Family for anti-ACVR2A antibody

[Epigenetics](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	518828583
NCBI GeneID	92
NCBI Accession #	NP_001265508.1
NCBI GenBank Nucleotide #	NM_001278579.1
Molecular Weight	45,713 Da
NCBI Official Full Name	activin receptor type-2A isoform 1
NCBI Official Synonym Full Names	activin A receptor, type IIA
NCBI Official Symbol	ACVR2A
NCBI Official Synonym Symbols	ACVR2; ACTRII
NCBI Protein Information	activin receptor type-2A
UniProt Protein Name	Activin receptor type-2A
UniProt Gene Name	ACVR2A
UniProt Synonym Gene Names	ACVR2; ACTR-IIA; ACTRIIA
UniProt Entry Name	AVR2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.