

Rabbit TRAF3 Polyclonal Antibody | anti-TRAF3 antibody

TRAF3 Polyclonal Antibody

CATALOG # AAA169244	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

TRAF3; Polyclonal Antibody; TRAF3 Polyclonal Antibody; CAP1; LAP1; CAP-1; CRAF1; IIAE5; CD40bp; anti-TRAF3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1.3mg/mL (varies by lot)
Applicable Applications for anti-TRAF3 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Synthetic peptide of human TRAF3
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human lung cancer using TRAF3 Polyclonal Antibody at dilution of 1:40



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human ovarian cancer using TRAF3 Polyclonal Antibody at dilution of 1:40



WB (Western Blot)

Western Blot analysis of Human fetal kidney tissue using TRAF3 Polyclonal Antibody at dilution of 1:500

ADDITIONAL INFORMATION

Related Product Information for anti-TRAF3 antibody

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from, members of the TNF receptor (TNFR) superfamily. This protein participates in the signal transduction of CD40, a TNFR family member important for the activation of the immune response. This protein is found to be a critical component of the lymphotoxin-beta receptor (LTbetaR) signaling complex, which induces NF-kappaB activation and cell death initiated by LTbeta ligation. Epstein-Barr virus encoded latent infection membrane protein-1 (LMP1) can interact with this and several other members of the TRAF family, which may be essential for the oncogenic effects of LMP1. Several alternatively spliced transcript variants encoding three distinct isoforms have been reported.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/traf3-antibody-immunohistochemistry-western-blot-elisa/169244>