

Rabbit LCN2 Polyclonal Antibody | anti-LCN2 antibody

LCN2 Antibody

CATALOG # AAA169313	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

LCN2; Polyclonal Antibody; LCN2 Antibody; 24p3; MSFI; NGAL; anti-LCN2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1mg/mL (varies by lot)
Sequence Length	198
Applicable Applications for anti-LCN2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human LCN2
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES



Application Data



WB (Western Blot)

Western blot analysis of extracts of mouse liver cancer tissue lysate, using LCN2 antibody. / Immunohistochemistry of paraffin-embedded human breast cancer tissue using LCN2 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-LCN2 antibody	Iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and renal development. Binds iron through association with 2, 5-dihydroxybenzoic acid (2, 5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin, and delivers or removes iron from the cell, depending on the context. Iron-bound form (holo-24p3) is internalized following binding to the SLC22A17 (24p3R) receptor, leading to release of iron and subsequent increase of intracellular iron concentration. In contrast, association of the iron-free form (apo-24p3) with the SLC22A17 (24p3R) receptor is followed by association with an intracellular siderophore, iron chelation and iron transfer to the extracellular medium, thereby reducing intracellular iron concentration. Involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis, while iron-free form decreases intracellular iron levels, inducing expression of the proapoptotic protein BCL2L11/BIM, resulting in apoptosis.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	49457137
NCBI GeneID	3934
UniProt Accession #	P80188; P30150; Q5SYV9; Q5SYW0; Q6FGL5; Q92683; A6NII8; B4DWV4; B7ZAA2
Molecular Weight	22,457 Da
NCBI Official Full Name	LCN2, partial
NCBI Official Synonym Full Names	lipocalin 2
NCBI Official Symbol	LCN2
NCBI Official Synonym Symbols	24p3; MSFI; NGAL
NCBI Protein Information	neutrophil gelatinase-associated lipocalin; 25 kDa alpha-2-microglobulin-related subunit of MMP-9; lipocalin-2; migration-stimulating factor inhibitor; oncogene 24p3; p25; siderocalin LCN2
UniProt Protein Name	Neutrophil gelatinase-associated lipocalin
UniProt Gene Name	LCN2
UniProt Synonym Gene Names	HNL; NGAL; NGAL
UniProt Entry Name	NGAL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/lcn2-antibody-immunohistochemistry-western-blot-elisa/169313>