

Rabbit CCL17 Polyclonal Antibody | anti-CCL17 antibody

CCL17 Polyclonal Antibody

CATALOG # AAA169645	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

CCL17; Polyclonal Antibody; CCL17 Polyclonal Antibody; TARC; ABCD-2; SCYA17; A-152E5.3; anti-CCL17 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2.6mg/mL (varies by lot)
Sequence Length	94
Applicable Applications for anti-CCL17 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Synthetic peptide of human CCL17
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.


IMAGES



IHC (Immunohistochemisry)
Immunohistochemistry of paraffin-embedded Human gastric cancer using CCL17 Polyclonal Antibody at dilution of 1:70



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded Human colon cancer using CCL17 Polyclonal Antibody at dilution of 1:70



WB (Western Blot)

Western Blot analysis of Mouse lung and thymus tissue, Human fetal muscle tissue using CCL17 Polyclonal Antibody at dilution of 1:1300

ADDITIONAL INFORMATION

Related Product Information for anti-CCL17 antibody

This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506829
NCBI GeneID	6361
NCBI Accession #	NP_002978.1
NCBI GenBank Nucleotide #	NM_002987.2
Molecular Weight	10,507 Da
NCBI Official Full Name	C-C motif chemokine 17
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 17
NCBI Official Symbol	CCL17
NCBI Official Synonym Symbols	TARC; ABCD-2; SCYA17; A-152E5.3
NCBI Protein Information	C-C motif chemokine 17; CC chemokine TARC; T cell-directed CC chemokine; small inducible cytokine subfamily A (Cys-Cys), member 17; small-inducible cytokine A17; thymus and activation-regulated chemokine
UniProt Protein Name	C-C motif chemokine 17
UniProt Gene Name	CCL17
UniProt Synonym Gene Names	SCYA17; TARC
UniProt Entry Name	CCL17_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ccl17-antibody-immunohistochemistry-western-blot-elisa/169645>