

Rabbit PLAU Polyclonal Antibody | anti-PLAU antibody

PLAU Polyclonal Antibody

CATALOG # AAA170130	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

PLAU; Polyclonal Antibody; PLAU Polyclonal Antibody; ATF; QPD; UPA; URK; u-PA; BDPLT5; anti-PLAU antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	0.3mg/mL (varies by lot)
Sequence Length	244
Applicable Applications for anti-PLAU antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant protein of human PLAU
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using PLAU Polyclonal Antibody at dilution of 1:10



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon cancer tissue using PLAU Polyclonal Antibody at dilution of 1:10

ADDITIONAL INFORMATION

Related Product Information for anti-PLAU antibody	This gene encodes a serine protease involved in degradation of the extracellular matrix and possibly tumor cell migration and proliferation. A specific polymorphism in this gene may be associated with late-onset Alzheimer's disease and also with decreased affinity for fibrin-binding. This protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-Ile bond to form a two-chain derivative in which a single disulfide bond connects the amino-terminal A-chain to the catalytically active, carboxy-terminal B-chain. This two-chain derivative is also called HMW-uPA (high molecular weight uPA). HMW-uPA can be further processed into LMW-uPA (low molecular weight uPA) by cleavage of chain A into a short chain A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA receptor. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-PLAU antibody	Epigenetics

NCBI AND UNIPROT INFORMATION

NCBI GI #	332743784
Molecular Weight	24,535 Da
NCBI Official Full Name	PlaU
UniProt Protein Name	PlaU
UniProt Gene Name	plaU
UniProt Entry Name	Q2I772_9ACTO

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/plau-antibody-immunohistochemistry-elisa/170130>