

Rabbit RPS27 Polyclonal Antibody | anti-RPS27 antibody

RPS27 Polyclonal Antibody

CATALOG # AAA170685	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

RPS27; Polyclonal Antibody; RPS27 Polyclonal Antibody; S27; MPS1; MPS-1; anti-RPS27 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1.2mg/mL (varies by lot)
Sequence Length	28
Applicable Applications for anti-RPS27 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human RPS27
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human liver cancer using RPS27 Polyclonal Antibody at dilution of 1:25



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human thyroid cancer using RPS27 Polyclonal Antibody at dilution of 1:25



WB (Western Blot)

Western Blot analysis of Jurkat, K562 and Raji cell using RPS27 Polyclonal Antibody at dilution of 1:300

ADDITIONAL INFORMATION

Related Product Information for anti-RPS27 antibody

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 40S subunit. The protein belongs to the S27E family of ribosomal proteins. It contains a C4-type zinc finger domain that can bind to zinc. The encoded protein has been shown to be able to bind to nucleic acid. It is located in the cytoplasm as a ribosomal component, but it has also been detected in the nucleus. Studies in rat indicate that ribosomal protein S27 is located near ribosomal protein S18 in the 40S subunit and is covalently linked to translation initiation factor eIF3. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

NCBI AND UNIPROT INFORMATION

NCBI GI #	728674869
NCBI Official Full Name	rps27

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/rps27-antibody-immunohistochemistry-western-blot-elisa/170685>