

## Human AICAR (AICA Riboside) CLIA Kit | AICAR clia kit

Human AICAR (AICA Riboside) CLIA Kit

CATALOG #  
**AAA171284**

REACTIVITY  
**Human**


### PRODUCT SYNONYMS

AICAR (AICA Riboside); Human AICAR (AICA Riboside) CLIA Kit; AICAR clia kit

### PRODUCT SPECIFICATIONS

|                 |                                                                                                                                              |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------|
| Reactivity      | Human                                                                                                                                        |
| Specificity     | This kit recognizes natural and recombinant AICAR. No significant cross-reactivity or interference between AICAR and analogues was observed. |
| Assay Type      | Sandwich                                                                                                                                     |
| Detection Range | 7.813-500pg/mL                                                                                                                               |
| Sensitivity     | Min: 4.688pg/mL; Max: 500pg/mL                                                                                                               |

### IMAGES



Standard Curve (Sample)

### ADDITIONAL INFORMATION

#### Related Product Information for AICAR clia kit

**Intended Uses:** This CLIA kit applies to the in vitro quantitative determination of AICAR concentrations in serum, plasma and other biological fluids.

**Principle of the Assay:** This kit uses Sandwich-CLIA as the method. The micro CLIA plate provided in this kit has been pre-coated with an antibody specific to AICAR. Standards or samples are added to the appropriate micro CLIA plate wells and combined with the specific antibody. Then a biotinylated detection antibody specific for AICAR and Avidin-Horseradish Peroxidase (HRP) conjugate is added to each micro plate well successively and incubated. Free components are washed away. The substrate solution is added to each well. Only those wells that contain AICAR, biotinylated detection antibody and Avidin-HRP conjugate will appear fluorescence. The Relative light unit (RLU) value is measured spectrophotometrically by the Chemiluminescence immunoassay analyzer. The RLU value is positively associated with the concentration of AICAR. You can calculate the concentration of AICAR in the samples by comparing the RLU value of the samples to the standard curve.

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

