

Rabbit DLAT Polyclonal Antibody | anti-DLAT antibody

DLAT Polyclonal Antibody

CATALOG # AAA173145	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

DLAT; Polyclonal Antibody; DLAT Polyclonal Antibody; DLTA; PDCE2; PDC-E2; anti-DLAT antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Concentration	2.3mg/ml (varies by lot)
Sequence Length	553
Applicable Applications for anti-DLAT antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Fusion protein of human DLAT
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-166899 / sc-365276
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human prostate cancer using DLAT Polyclonal Antibody at dilution of 1/40


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human breast cancer using DLAT Polyclonal Antibody at dilution of 1/40

ADDITIONAL INFORMATION

Related Product Information for anti-DLAT antibody	This gene encodes component E2 of the multi-enzyme pyruvate dehydrogenase complex (PDC). PDC resides in the inner mitochondrial membrane and catalyzes the conversion of pyruvate to acetyl coenzyme A. The protein product of this gene, dihydrolipoamide acetyltransferase, accepts acetyl groups formed by the oxidative decarboxylation of pyruvate and transfers them to coenzyme A. Dihydrolipoamide acetyltransferase is the antigen for antimitochondrial antibodies. These autoantibodies are present in nearly 95% of patients with the autoimmune liver disease primary biliary cirrhosis (PBC). In PBC, activated T lymphocytes attack and destroy epithelial cells in the bile duct where this protein is abnormally distributed and overexpressed. PBC eventually leads to cirrhosis and liver failure. Mutations in this gene are also a cause of pyruvate dehydrogenase E2 deficiency which causes primary lactic acidosis in infancy and early childhood.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	755919262
Molecular Weight	69 kDa
NCBI Official Full Name	DlaT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/dlat-antibody-immunohistochemistry-western-blot-elisa/173145>