

CD171 / N-CAML1 / L1CAM recombinant protein

Recombinant Human CD171/L1CAM Protein (His Tag)

CATALOG #
AAA173574

HOST
HEK293 Cells


PRODUCT SYNONYMS

CD171 / N-CAML1 / L1CAM; Recombinant Human CD171/L1CAM Protein (His Tag); CAML1; CD171; HSAS; HSAS1; MASA; MIC5; N-CAM-L1; N-CAML1; NCAML1; S10; SPG1; CD171 / N-CAML1 / L1CAM recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	> 85 % as determined by reducing SDS-PAGE.
Form/Format	Lyophilized from sterile PBS, pH 7.4, 5% Trehalose, 5% Mannitol, 0.01% Tween 80.
Sequence	Met1-Glu11 20
Species	Human
Tag	C-His
Endotoxin	< 1.0 EU per ug as determined by the LAL method.
Reconstitution	It is recommended that sterile water (400uL) be added to the vial to prepare a stock solution of 0.25mg/mL. Concentration is measured by UV-Vis.
Preparation and Storage	Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80 degree C. Reconstituted protein solution can be stored at 4-8 degree C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 degree C for 3 months. Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for CD171 / N-CAML1 / L1CAM recombinant protein	Background: L1 cell adhesion molecule (L1CAM), also designated as CD171, is a cell adhesion receptor of the immunoglobulin superfamily, known for its roles in nerve cell function. While originally believed to be present only in brain cells, in recent years L1-CAM has been detected in other tissues, and in a variety of cancer cells, including some common type s of human cancer. L1CAM interacts with a variety of ligands including axonin-1, CD9, neurocan and integrins, and it has been revealed that the RGD motif in the sixth Ig domain of L1CAM is a binding site for integrins, thus important for nuclear signaling. Disruption of L1CAM function causes three X-linked neurological syndromes, i.e. hydrocephalus, MASA syndrome (mental retardation, aphasia, shuffling gait and adducted thumbs) and spastic paraplegia syndrome. Overexpression of L1CAM in normal and cancer cells increased motility, enhanced growth rate and promoted cell transformation and tumorigenicity. Recent work has identified L1CAM (CD171) as a novel marker for human carcinoma progression, and a candidate for anti-cancer therapy. Description: A DNA sequence encoding the extracellular domain of human CD171 (NP_000416.1) (Met 1-Glu 1120) was fused with a polyhistidine tag at the C-terminus.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557707
NCBI GeneID	3897
NCBI Accession #	NP_000416.1
NCBI GenBank Nucleotide #	NM_000425.4
Molecular Weight	Mol Mass: 125 kDa AP Mol Mass: 160-200 kDa
NCBI Official Full Name	neural cell adhesion molecule L1 isoform 1
NCBI Official Synonym Full Names	L1 cell adhesion molecule
NCBI Official Symbol	L1CAM
NCBI Official Synonym Symbols	S10; HSAS; MASA; MIC5; SPG1; CAML1; CD171; HSAS1; N-CAML1; NCAM-L1; N-CAM-L1
NCBI Protein Information	neural cell adhesion molecule L1
UniProt Protein Name	Neural cell adhesion molecule L1
UniProt Gene Name	L1CAM
UniProt Synonym Gene Names	CAML1; MIC5; N-CAM-L1; NCAM-L1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/l1cam-recombinant-protein-hek293/173574>