

Rabbit TGFbeta2 Polyclonal Antibody | anti-TGFbeta2 antibody

TGFbeta2 Polyclonal Antibody

CATALOG # AAA173922	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

TGFbeta2; Polyclonal Antibody; TGFbeta2 Polyclonal Antibody; Transforming growth factor beta-2 (TGF-beta-2) (BSC-1 cell growth inhibitor) (Cetermin) (Glioblastoma-derived T-cell suppressor factor) (G-TSF) (Polygerin); anti-TGFbeta2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/mL (varies by lot)
Sequence Length	442
Applicable Applications for anti-TGFbeta2 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Synthetic peptide from human protein
Buffer	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.


IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human kidney tissue using TGFbeta2 Polyclonal Antibody at dilution of 1:200.



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human kidney tissue using TGFbeta2 Polyclonal Antibody at dilution of 1:200.



WB (Western Blot)

Western Blot analysis of Mouse kidney, 293T, Mouse lung using TGFbeta2 Polyclonal Antibody at dilution of 1:1000.

ADDITIONAL INFORMATION

Related Product Information for anti-TGFbeta2 antibody

This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types by transducing their signal through combinations of transmembrane type I and type II receptors (TGFB1 and TGFB2) and their downstream effectors, the SMAD proteins. Disruption of the TGFB/SMAD pathway has been implicated in a variety of human cancers. The encoded protein is secreted and has suppressive effects of interleukin-2 dependent T-cell growth. Translocation t(1;7)(q41;p21) between this gene and HDAC9 is associated with Peters' anomaly, a congenital defect of the anterior chamber of the eye. The knockout mice lacking this gene show perinatal mortality and a wide range of developmental, including cardiac, defects. Alternatively spliced transcript variants encoding different isoforms have been identified.

NCBI AND UNIPROT INFORMATION

NCBI GI #	208022653
NCBI GeneID	7042
NCBI Accession #	NP_001129071.1
NCBI GenBank Nucleotide #	NM_001135599.3
UniProt Accession #	P61812; P08112; Q15579; Q15581; Q4VAV9; B4DKC5
Molecular Weight	47+12kDa
NCBI Official Full Name	transforming growth factor beta-2 isoform 1
NCBI Official Synonym Full Names	transforming growth factor beta 2
NCBI Official Symbol	TGFB2
NCBI Official Synonym Symbols	LDS4; G-TSF; TGF-beta2
NCBI Protein Information	transforming growth factor beta-2
UniProt Protein Name	Transforming growth factor beta-2
UniProt Gene Name	TGFB2
UniProt Synonym Gene Names	TGF-beta-2; G-TSF; LAP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/tgfbeta2-antibody-immunohistochemistry-western-blot-elisa/173922>