

Rabbit anti-Human, Mouse MIP-3alpha Polyclonal Antibody | anti-MIP-3alpha antibody

MIP-3alpha Polyclonal Antibody

CATALOG # AAA173928	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS ELISA, Immunohistochemistry
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

MIP-3alpha; Polyclonal Antibody; MIP-3alpha Polyclonal Antibody; C-C motif chemokine 20 (Beta-chemokine exodus-1) (CC chemokine LARC) (Liver and activation-regulated chemokine) (Macrophage inflammatory protein 3 alpha) (MIP-3-alpha) (Small-inducible cytokine A20) [Cleaved into: CCL20(1-67); CCL20(1-64); CCL20(2-70)]; anti-MIP-3alpha antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/mL (varies by lot)
Sequence Length	95
Applicable Applications for anti-MIP-3alpha antibody	ELISA, IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein
Buffer	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human pancreas tissue using MIP-3alpha Polyclonal Antibody at dilution of 1:200.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human liver tissue using MIP-3alpha Polyclonal Antibody at dilution of 1:200.

ADDITIONAL INFORMATION

Related Product Information for anti-MIP-3alpha antibody	CCL20 (C-C Motif Chemokine Ligand 20) is a Protein Coding gene. Diseases associated with CCL20 include Bejel and Rheumatoid Arthritis. Among its related pathways are Bacterial infections in CF airways and Peptide ligand-binding receptors. GO annotations related to this gene include cytokine activity and chemokine activity. This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene displays chemotactic activity for lymphocytes and can repress proliferation of myeloid progenitors. Two transcript variants encoding different isoforms have been found for this gene.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	194239688
NCBI GeneID	6364
NCBI Accession #	NP_001123518.1
NCBI GenBank Nucleotide #	NM_001130046.1
UniProt Accession #	P78556; Q53S51; Q99664
Molecular Weight	10,691 Da
NCBI Official Full Name	C-C motif chemokine 20 isoform 2
NCBI Official Synonym Full Names	C-C motif chemokine ligand 20
NCBI Official Symbol	CCL20
NCBI Official Synonym Symbols	CKb4; LARC; ST38; MIP3A; Exodus; MIP-3a; SCYA20; MIP-3-alpha
NCBI Protein Information	C-C motif chemokine 20
UniProt Protein Name	C-C motif chemokine 20
UniProt Gene Name	CCL20
UniProt Synonym Gene Names	LARC; MIP3A; SCYA20; MIP-3-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/mip-3alpha-antibody-immunohistochemistry-elisa/173928>