

Rabbit anti-Human IL-12B p40 Polyclonal Antibody | anti-IL-12B p40 antibody

IL-12B p40 Polyclonal Antibody

CATALOG # AAA173945	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

IL-12B p40; Polyclonal Antibody; IL-12B p40 Polyclonal Antibody; Interleukin-12 subunit beta (IL-12B) (Cytotoxic lymphocyte maturation factor 40 kDa subunit) (CLMF p40) (IL-12 subunit p40) (NK cell stimulatory factor chain 2) (NKSF2); anti-IL-12B p40 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/mL (varies by lot)
Sequence Length	328
Applicable Applications for anti-IL-12B p40 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Synthetic peptide from human protein
Buffer	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon tissue using IL-12B p40 Polyclonal Antibody at dilution of 1:200.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon tissue using IL-12B p40 Polyclonal Antibody at dilution of 1:200.



WB (Western Blot)

Western Blot analysis of Mouse kidney using IL-12B p40 Polyclonal Antibody at dilution of 1:1000.

ADDITIONAL INFORMATION

Related Product Information for anti-IL-12B p40 antibody

Interleukin-12 (IL-12), is a cytokine that is secreted by activated phagocytes and dendritic cells and that induces interferon-gamma production by natural-killer and T lymphocytes. IL-12 is a 75 kDa heterodimer composed of a 35 kDa subunit (IL-12A p35) and a 40 kDa subunit (IL-12B p40) that is secreted by a wide variety of antigen presenting cells (APCs), including phagocytes, B cells and Langerhans cells. IL-12B has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen. Overexpression of IL-12B was observed in the central nervous system of patients with multiple sclerosis (MS), suggesting a role of this cytokine in the pathogenesis of the disease. The promoter polymorphism of IL-12B has been reported to be associated with the severity of atopic and non-atopic asthma in children.

NCBI AND UNIPROT INFORMATION

NCBI GI #	24497438
NCBI GeneID	3593
NCBI Accession #	NP_002178.2
NCBI GenBank Nucleotide #	NM_002187.2
UniProt Accession #	P29460
Molecular Weight	40kDa
NCBI Official Full Name	interleukin-12 subunit beta
NCBI Official Synonym Full Names	interleukin 12B
NCBI Official Symbol	IL12B
NCBI Official Synonym Symbols	CLMF; NKSF; CLMF2; IMD28; IMD29; NKSF2; IL-12B
NCBI Protein Information	interleukin-12 subunit beta
UniProt Protein Name	Interleukin-12 subunit beta
UniProt Gene Name	IL12B
UniProt Synonym Gene Names	NKSF2; IL-12B; CLMF p40; NKSF2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/il-12b-p40-antibody-immunohistochemistry-western-blot-elisa/173945>