

Activin Receptor 2A/Activin RIIA/ACVR2A Recombinant Protein | ACVR2A recombinant protein

Recombinant Human Activin Receptor 2A/Activin RIIA/ACVR2A Protein (C-Fc-6His)

CATALOG # AAA174004	HOST Human Cells
-------------------------------	----------------------------


PRODUCT SYNONYMS

Activin Receptor 2A/Activin RIIA/ACVR2A; Recombinant Human Activin Receptor 2A/Activin RIIA/ACVR2A Protein (C-Fc-6His); Activin Receptor Type-2A; Activin Receptor Type IIA; ACTR-IIA; ACTRIIA; ACVR2A; ACVR2; ACVR2A recombinant protein

PRODUCT SPECIFICATIONS

Host	Human Cells
Purity/Purification	>95% as determined by reducing SDS-PAGE.
Form/Format	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Sequence Positions	Ala20-Pro134
Sequence Length	513
Species	Human
Endotoxin	<1.0 EU per µg as determined by LAL test.
Fusion Tag	C-Fc-6His
Bio Activity	Immobilized Human INHBC-His at 10µg/mL (100 µl/well) can bind Human ACVR2A-Fc-6His. The ED50 of Human ACVR2A-Fc-6His is 2-8 µg/mL
Reconstitution	We recommend that the vial be centrifuged to opening to bring the contents to the bottom. Do not mix by vortex. It is recommended that sterile water be added to the vial to prepare a stock solution of 0.25 mg/mL. Concentration is measured by UV-Vis.
Preparation and Storage	Lyophilized protein should be stored at < -20 degree C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7 degree C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 degree C for 3 months.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for ACVR2A recombinant protein	Activin Receptor Type-2A is a protein that in humans is encoded by the ACVR2A gene. ACVR2A is an activin type 2 receptor. This gene encodes activin A type II receptor. Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	518828583
NCBI GeneID	92
NCBI Accession #	NP_001265508.1
NCBI GenBank Nucleotide #	NM_001278579.1
UniProt Accession #	P27037; Q53TH4; Q6NWW2; Q92474; B2RAB8; B4DWQ2; D3DP85
Molecular Weight	Molecular Mass: 41.2 kDa Actual Protein Molecular Mass: 36 kDa
NCBI Official Full Name	activin receptor type-2A isoform 1
NCBI Official Synonym Full Names	activin A receptor type 2A
NCBI Official Symbol	ACVR2A
NCBI Official Synonym Symbols	ACVR2; ACTRII
NCBI Protein Information	activin receptor type-2A
UniProt Protein Name	Activin receptor type-2A
UniProt Gene Name	ACVR2A
UniProt Synonym Gene Names	ACVR2; ACTR-IIA; ACTRIIA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/acvr2a-recombinant-protein-human/174004>