

Mesothelin Recombinant Protein | MSLN recombinant protein

Recombinant Human Mesothelin Protein

CATALOG # AAA174585	HOST Human Cells
-------------------------------	----------------------------


PRODUCT SYNONYMS

Mesothelin; Recombinant Human Mesothelin Protein; Megakaryocyte potentiating factor; mesothelin; Pre-pro-megakaryocyte-potentiating factor; soluble MPF mesothelin related protein; CAK1; MPF; MSLN; SMR; CAK1 antigen; MSLN recombinant protein

PRODUCT SPECIFICATIONS

Host	Human Cells
Purity/Purification	>95% as determined by reducing SDS-PAGE.
Form/Format	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Sequence Length	622
Species	Human
Endotoxin	<1.0 EU per µg as determined by LAL test.
Fragment	Glu296-Ser598
Fusion Tag	C-6His
Reconstitution	We recommend that the vial be centrifuged to opening to bring the contents to the bottom. Do not mix by vortex. It is recommended that sterile water be added to the vial to prepare a stock solution of 0.25 mg/mL. Concentration is measured by UV-Vis.
Preparation and Storage	Lyophilized protein should be stored at < -20 degree C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7 degree C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 degree C for 3 months.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for MSLN recombinant protein	Protein Construction: Recombinant Human Mesothelin is produced by our Mammalian expression system and the target gene encoding Glu296-Ser598 is expressed with a 6His tag at the C-terminus. Background: Mesothelin is a cell surface glycoprotein whose expression is limited to mesothelial cells of the serosa (pleura, pericardium, and peritoneum) and epithelial cells of the trachea, tonsils, fallopian tube, and kidneys. Mesothelin plays an important role in cell survival, proliferation, migration, invasion, tumor progression, and resistance to chemotherapy. The overexpression of mesothelin can activate NF-kappaB and signal transducer and activator of transcription 3 (Stat3), inhibit apoptotic signaling and TNF-alpha-induced apoptosis, and accelerate the G1-S transition. Mesothelin is also found overexpressed in various cancers, including malignant mesothelioma, pancreatic or ovarian carcinoma, sarcomas and in some gastrointestinal or pulmonary carcinomas. As a result of its limited expression in normal tissues, mesothelin has been reported as an ideal tumor-associated marker for the development of targeted therapy.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	293651530
NCBI GeneID	10232
NCBI Accession #	NP_001170826.1
NCBI GenBank Nucleotide #	NM_001177355.2
UniProt Accession #	Q13421; Q14859; Q4VQD5; Q96GR6; Q96KJ5; Q9BR17; Q9BTR2; Q9UCB2; Q9UK57; D3DU65
Molecular Weight	Mol_Mass: 34.9 kDa AP_Mol_Mass: 35-50 kDa
NCBI Official Full Name	mesothelin isoform 1 preproprotein
NCBI Official Synonym Full Names	mesothelin
NCBI Official Symbol	MSLN
NCBI Official Synonym Symbols	MPF; SMRP
NCBI Protein Information	mesothelin
UniProt Protein Name	Mesothelin
UniProt Gene Name	MSLN
UniProt Synonym Gene Names	MPF; MPF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.