

Rabbit anti-Human COMT Polyclonal Antibody | anti-COMT antibody

COMT Polyclonal Antibody

CATALOG # AAA175283	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

COMT; Polyclonal Antibody; COMT Polyclonal Antibody; Catechol O methyltransferase; Catechol O-methyltransferase; EC 2.1.1.6; anti-COMT antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Form/Format	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4
Concentration	1.02mg/mL (varies by lot)
Sequence Length	271
Applicable Applications for anti-COMT antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Fusion protein of human COMT
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using COMT Polyclonal Antibody at dilution of 1:30(×200)



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using COMT Polyclonal Antibody at dilution of 1:30(×200)



WB (Western Blot)

Western blot analysis of Hela cell lysate using COMT Polyclonal Antibody at dilution of 1:450

ADDITIONAL INFORMATION

Related Product Information for anti-COMT antibody

Catechol-O-methyltransferase catalyzes the transfer of a methyl group from S-adenosylmethionine to catecholamines, including the neurotransmitters dopamine, epinephrine, and norepinephrine. This O-methylation results in one of the major degradative pathways of the catecholamine transmitters. In addition to its role in the metabolism of endogenous substances, COMT is important in the metabolism of catechol drugs used in the treatment of hypertension, asthma, and Parkinson disease. COMT is found in two forms in tissues, a soluble form (S-COMT) and a membrane-bound form (MB-COMT). The differences between S-COMT and MB-COMT reside within the N-termini. Several transcript variants are formed through the use of alternative translation initiation sites and promoters.

Product Categories/Family for anti-COMT antibody

[Cancer](#); [Metabolism](#); [Neuroscience](#); [Signal transduction](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502969
NCBI GeneID	1312
NCBI Accession #	NP_000745.1; BC011935
NCBI GenBank Nucleotide #	NP_000745.1
UniProt Accession #	P21964
Molecular Weight	30 kDa
NCBI Official Full Name	catechol O-methyltransferase isoform MB-COMT
UniProt Protein Name	Catechol O-methyltransferase
UniProt Gene Name	COMT
UniProt Entry Name	COMT_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/comt-antibody-immunohistochemistry-western-blot-elisa/175283>