

Rabbit anti-Human, Mouse EWSR1 Polyclonal Antibody | anti-EWSR1 antibody

EWSR1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA175487	Rabbit	Human, Mouse	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

EWSR1; Polyclonal Antibody; EWSR1 Polyclonal Antibody; bK984G1.4; BK984G1.4 Ewing sarcoma breakpoint region 1 protein; Ewing sarcoma breakpoint region 1; Ewing sarcoma breakpoint region 1 protein; Ewings sarcoma EWS Fli1 type 1 oncogene; EWS; EWS oncogene; EWS RNA binding protein 1; EWSR 1; Ewsr1; EWSR1 protein; RNA binding protein EWS; RNA-binding protein EWS; anti-EWSR1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Form/Format	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4
Concentration	1.26mg/mL (varies by lot)
Sequence Length	655
Applicable Applications for anti-EWSR1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Fusion protein of human EWSR1
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human lung cancer tissue using EWSR1 Polyclonal Antibody at dilution of 1:70(×200)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using EWSR1 Polyclonal Antibody at dilution of 1:70(×200)



WB (Western Blot)
 Western blot analysis of 293T and HepG2 cell lysates using EWSR1 Polyclonal Antibody at dilution of 1:1000

ADDITIONAL INFORMATION

Related Product Information for anti-EWSR1 antibody	This gene encodes a multifunctional protein that is involved in various cellular processes, including gene expression, cell signaling, and RNA processing and transport. The protein includes an N-terminal transcriptional activation domain and a C-terminal RNA-binding domain. Chromosomal translocations between this gene and various genes encoding transcription factors result in the production of chimeric proteins that are involved in tumorigenesis. These chimeric proteins usually consist of the N-terminal transcriptional activation domain of this protein fused to the C-terminal DNA-binding domain of the transcription factor protein. Mutations in this gene, specifically a t(11;22)(q24;q12) translocation, are known to cause Ewing sarcoma as well as neuroectodermal and various other tumors. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1 and 14.
Product Categories/Family for anti-EWSR1 antibody	Cancer ; Epigenetics and Nuclear Signaling ; Tags and Cell Markers

NCBI AND UNIPROT INFORMATION

NCBI GI #	253970502
NCBI GeneID	2130
NCBI Accession #	NP_001156757.1; BC000527
NCBI GenBank Nucleotide #	NP_001156757.1
UniProt Accession #	Q01844
Molecular Weight	68 kDa
NCBI Official Full Name	RNA-binding protein EWS isoform 3
NCBI Official Synonym Full Names	EWS RNA binding protein 1
NCBI Official Symbol	EWSR1
NCBI Official Synonym Symbols	EWS; EWS-FLI1; bK984G1.4
NCBI Protein Information	RNA-binding protein EWS
UniProt Protein Name	RNA-binding protein EWS
UniProt Gene Name	EWSR1
UniProt Synonym Gene Names	EWS
UniProt Entry Name	EWS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ewsr1-antibody-immunohistochemistry-western-blot-elisa/175487>