

Rabbit Androgen Receptor Polyclonal Antibody | anti-AR antibody

Androgen Receptor Polyclonal Antibody

CATALOG # AAA178712	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
-------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

Androgen Receptor; Polyclonal Antibody; Androgen Receptor Polyclonal Antibody; AIS; AR8; DHTR; HUMARA; HYSP1; KD; NR3C4; SBMA; SMAX1; TFM; AR; anti-AR antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-AR antibody	WB (Western Blot)
Immunogen	A synthetic peptide of human Androgen Receptor
Research Area	Cancer, Epigenetics and Nuclear Signaling, Developmental Biology, Signal Transduction
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES


WB (Western Blot)

Western blot analysis of extracts of Mouse testis using Androgen Receptor Polyclonal Antibody at 1:500 dilution.


WB (Western Blot)

Western blot analysis of extracts of Rat testis using Androgen Receptor Polyclonal Antibody at 1:500 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-AR antibody	The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two alternatively spliced variants encoding distinct isoforms have been described.
Product Categories/Family for anti-AR antibody	Antibodies; Primary Antibodies; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	21322252
NCBI GeneID	367
NCBI Accession #	NP_000035.2
NCBI GenBank Nucleotide #	NM_000044.3
UniProt Accession #	P10275; Q9UD95; A2RUN2; B1AKD7
Molecular Weight	98,989 Da
NCBI Official Full Name	androgen receptor isoform 1
NCBI Official Synonym Full Names	androgen receptor
NCBI Official Symbol	AR
NCBI Official Synonym Symbols	KD; AIS; TFM; DHTR; SBMA; HYS P1; NR3C4; SMAX1; HUMARA
NCBI Protein Information	androgen receptor; dihydrotestosterone receptor; androgen nuclear receptor variant 2; nuclear receptor subfamily 3 group C member 4
UniProt Protein Name	Androgen receptor
UniProt Gene Name	AR
UniProt Synonym Gene Names	DHTR; NR3C4
UniProt Entry Name	ANDR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/androgen-receptor-antibody-western-blot-elisa/178712>