

Rabbit 53BP1 Polyclonal Antibody | anti-53BP1 antibody

53BP1 Polyclonal Antibody

CATALOG # AAA178739	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

53BP1; Polyclonal Antibody; 53BP1 Polyclonal Antibody; TP53BP1; TDRD30; TP53; p202; p53BP1; anti-53BP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-53BP1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human 53BP1
Research Area	Cancer, Epigenetics and Nuclear Signaling
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human brain using 53BP1 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of HeLa cells using 53BP1 Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-53BP1 antibody	Double-strand break (DSB repair protein involved in response to DNA damage, telomere dynamics and class-switch recombination (CSR during antibody genesis. Plays a key role in the repair of double-strand DNA breaks (DSBs in response to DNA damage by promoting non-homologous end joining (NHEJ-mediated repair of DSBs and specifically counteracting the function of the homologous recombination (HR repair protein BRCA1. In response to DSBs, phosphorylation by ATM promotes interaction with RIF1 and dissociation from NUDT16L1/TIRR, leading to recruitment to DSBs sites. Recruited to DSBs sites by recognizing and binding histone H2A monoubiquitinated at 'Lys-15' (H2AK15Ub and histone H4 dimethylated at 'Lys-20' (H4K20me2, two histone marks that are present at DSBs sites. Required for immunoglobulin class-switch recombination (CSR during antibody genesis, a process that involves the generation of DNA DSBs. Participates in the repair and the orientation of the broken DNA ends during CSR (By similarity. In contrast, it is not required for classic NHEJ and V(DJ recombination (By similarity. Promotes NHEJ of dysfunctional telomeres via interaction with PAXIP1.
Product Categories/Family for anti-53BP1 antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	213972634
NCBI GeneID	7158
NCBI Accession #	NP_001135451.1
NCBI GenBank Nucleotide #	NM_001141979.1
UniProt Accession #	Q12888 ; Q2M1Z7 ; Q4LE46 ; Q5FWZ3 ; Q7Z3U4
Molecular Weight	213,574 Da
NCBI Official Full Name	tumor suppressor p53-binding protein 1 isoform 2
NCBI Official Synonym Full Names	tumor protein p53 binding protein 1
NCBI Official Symbol	TP53BP1
NCBI Official Synonym Symbols	p202; 53BP1
NCBI Protein Information	tumor suppressor p53-binding protein 1; p53BP1; tumor protein 53-binding protein, 1; tumor protein p53-binding protein, 1
UniProt Protein Name	Tumor suppressor p53-binding protein 1
UniProt Gene Name	TP53BP1
UniProt Synonym Gene Names	53BP1; p53-binding protein 1; p53BP1
UniProt Entry Name	TP53B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/53bp1-antibody-immunohistochemistry-western-blot-elisa/178739>