

Rabbit DBT Polyclonal Antibody | anti-DBT antibody

DBT Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA179260	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DBT; Polyclonal Antibody; DBT Polyclonal Antibody; E2; E2B; BCATE2; BCKADE2; BCKAD-E2; BCKDH-E2; BCOADC-E2; anti-DBT antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-DBT antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human DBT
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of PC-12 cells using DBT Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of A-431 cells using DBT Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse kidney using DBT Polyclonal Antibody at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines using DBT Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-DBT antibody	The branched-chain alpha-keto acid dehydrogenase complex (BCKD) is an inner-mitochondrial enzyme complex involved in the breakdown of the branched-chain amino acids isoleucine, leucine, and valine. The BCKD complex is thought to be composed of a core of 24 transacylase (E2) subunits, and associated decarboxylase (E1), dehydrogenase (E3), and regulatory subunits. This gene encodes the transacylase (E2) subunit. Mutations in this gene result in maple syrup urine disease, type 2. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.
Product Categories/Family for anti-DBT antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	400668
NCBI GeneID	1629
UniProt Accession #	P11182 ; Q5VVL8 ; B2R811
Molecular Weight	482
NCBI Official Full Name	Lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex, mitochondrial
NCBI Official Synonym Full Names	dihydrolipoamide branched chain transacylase E2
NCBI Official Symbol	DBT
NCBI Official Synonym Symbols	E2; E2B; BCATE2; BCKADE2; BCKAD-E2
NCBI Protein Information	lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex, mitochondrial; BCKAD E2 subunit; dihydrolipoyl transacylase; branched chain acyltransferase, E2 component; dihydrolipoylsine-residue (2-methylpropanoyl)transfer
UniProt Protein Name	Lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex, mitochondrial
UniProt Gene Name	DBT
UniProt Entry Name	ODB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/dbt-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/179260>