

Rabbit VCP Polyclonal Antibody | anti-VCP antibody

VCP Polyclonal Antibody

CATALOG # AAA179386	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

VCP; Polyclonal Antibody; VCP Polyclonal Antibody; ALS14; CDC48; CMT2Y; HEL-220; HEL-S-70; IBMPFD; IBMPFD1; TERA; p97; anti-VCP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.01% thiomersal, 50% glycerol, pH 7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-VCP antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human VCP
Research Area	Cancer, Metabolism, Neuroscience, Signal Transduction
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse stomach using VCP Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human esophageal using VCP Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat kidney using VCP Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines using VCP Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-VCP antibody	The protein encoded by this gene is member of a family that includes putative ATP-binding proteins involved in vesicle transport and fusion, 26S proteasome function, and assembly of peroxisomes. This protein, as a structural protein, is associated with clathrin, and heat-shock protein Hsc70, to form a complex. It has been implicated in a number of cellular events that are regulated during mitosis, including homotypic membrane fusion, spindle pole body function, and ubiquitin-dependent protein degradation.
Product Categories/Family for anti-VCP antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	6005942
NCBI GeneID	7415
NCBI Accession #	NP_009057.1
NCBI GenBank Nucleotide #	NM_007126.3
UniProt Accession #	P55072; Q0V924; Q2TAI5; Q969G7; Q9UCD5; B2R5T8
Molecular Weight	89,322 Da
NCBI Official Full Name	transitional endoplasmic reticulum ATPase
NCBI Official Synonym Full Names	valosin containing protein
NCBI Official Symbol	VCP
NCBI Official Synonym Symbols	p97; TERA; ALS14; IBMPFD
NCBI Protein Information	transitional endoplasmic reticulum ATPase; TER ATPase; yeast Cdc48p homolog; valosin-containing protein; 15S Mg(2+)-ATPase p97 subunit
UniProt Protein Name	Transitional endoplasmic reticulum ATPase
UniProt Gene Name	VCP
UniProt Synonym Gene Names	TER ATPase; VCP
UniProt Entry Name	TERA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

