

Rabbit TRAF7 Polyclonal Antibody | anti-TRAF7 antibody

TRAF7 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA179393	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TRAF7; Polyclonal Antibody; TRAF7 Polyclonal Antibody; RFWD1; RNF119; anti-TRAF7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300, 50% glycerol, pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TRAF7 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human TRAF7
Research Area	Cancer, Cell Biology, Immunology
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of PC-12 cells using TRAF7 Polyclonal Antibody at dilution of 1:50. Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using TRAF7 Polyclonal Antibody at dilution of 1:50. Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using TRAF7 Polyclonal Antibody at dilution of 1:50. Blue: DAPI for nuclear staining.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon carcinoma using TRAF7 Polyclonal Antibody at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



WB (Western Blot)

Western blot analysis of extracts of various cell lines using TRAF7 Polyclonal Antibody at 1:1000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-TRAF7 antibody	Tumor necrosis factor (TNF; see MIM 191160) receptor-associated factors, such as TRAF7, are signal transducers for members of the TNF receptor superfamily (see MIM 191190). TRAFs are composed of an N-terminal cysteine/histidine-rich region containing zinc RING and/or zinc finger motifs; a coiled-coil (leucine zipper) motif; and a homologous region that defines the TRAF family, the TRAF domain, which is involved in self-association and receptor binding.
Product Categories/Family for anti-TRAF7 antibody	Antibodies ; Primary Antibodies ; Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	54036486
NCBI GeneID	84231
UniProt Accession #	Q6Q0C0 ; Q9H073
Molecular Weight	670
NCBI Official Full Name	E3 ubiquitin-protein ligase TRAF7
NCBI Official Synonym Full Names	TNF receptor-associated factor 7, E3 ubiquitin protein ligase
NCBI Official Symbol	TRAF7
NCBI Official Synonym Symbols	RFWD1; RNF119

NCBI Protein Information	E3 ubiquitin-protein ligase TRAF7; RING finger protein 119; ring finger and WD repeat domain 1; RING finger and WD repeat-containing protein 1
UniProt Protein Name	E3 ubiquitin-protein ligase TRAF7
UniProt Gene Name	TRAF7
UniProt Synonym Gene Names	RFWD1; RNF119
UniProt Entry Name	TRAF7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/traf7-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/179393>