

Rabbit NF-kB p65 Polyclonal Antibody | anti-NF-kBp65 antibody

(KO Validated) NF-kB p65 Polyclonal Antibody

CATALOG # AAA179607	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

NF-kB p65; Polyclonal Antibody; (KO Validated) NF-kB p65 Polyclonal Antibody; RELA; NFKB3; p65; transcription factor p65; anti-NF-kBp65 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-NF-kBp65 antibody	IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant fusion protein of human NF-kB p65/RelA
Research Area	Cancer, Cell Biology, Cardiovascular, Epigenetics and Nuclear Signaling, Metabolism, Microbiology, Neuroscience, Signal Transduction
Preparation and Storage	Storage: -20 degree C/One year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of NIH/3T3 cells using [KO Validated] NF-kB p65/RelA Polyclonal Antibody at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.




IF (Immunofluorescence)

Immunofluorescence analysis of HepG2 cells using [KO Validated] NF-kB p65/RelA Polyclonal Antibody at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts from wild type(WT) and NF-kB p65/RelA knockout (KO) HeLa(KO) cells using NF-kB p65/RelA Polyclonal Antibody at 1:2000 dilution.



WB (Western Blot)

Western blot analysis of various lysates using NF-kB p65/RelA Polyclonal Antibody at 1:2000 dilution.

ADDITIONAL INFORMATION

Related Product Information for anti-NF-kBp65 antibody

NF-kappa-B is ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene.

Product Categories/Family for anti-NF-kBp65 antibody

[Antibodies](#); [Primary Antibodies](#); [Polyclonal Antibody](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	223468681
NCBI GeneID	5970
NCBI Accession #	NP_001138610.1
NCBI GenBank Nucleotide #	NM_001145138.1
UniProt Accession #	Q04206 ; Q6GTV1 ; Q6SLK1
Molecular Weight	60,219 Da
NCBI Official Full Name	transcription factor p65 isoform 2
NCBI Official Synonym Full Names	v-rel reticuloendotheliosis viral oncogene homolog A (avian)
NCBI Official Symbol	RELA
NCBI Official Synonym Symbols	p65; NFKB3
NCBI Protein Information	transcription factor p65; NF-kappa-B p65delta3; nuclear factor NF-kappa-B p65 subunit; nuclear factor of kappa light polypeptide gene enhancer in B-cells 3
UniProt Protein Name	Transcription factor p65
UniProt Gene Name	RELA
UniProt Synonym Gene Names	NFKB3
UniProt Entry Name	TF65_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

