

Rat platelet derived growth factor DD ELISA Kit | PDGF-DD elisa kit

Rat Platelet-derived growth factor DD, PDGF-DD ELISA Kit

CATALOG #
AAA18173

REACTIVITY
Rat


PRODUCT SYNONYMS

platelet derived growth factor DD; Rat Platelet-derived growth factor DD; PDGF-DD ELISA Kit; Rat Platelet-derived growth factor DD (PDGF-DD) ELISA kit; PDGF-DD elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of rat PDGF-DD. No significant cross-reactivity or interference between rat PDGF-DD and analogues was observed.
Sequence Length	300
Samples	Serum, plasma, tissue homogenates
Assay Type	Sandwich
Detection Range	0.156 ng/ml -10 ng/ml.
Sensitivity	The minimum detectable dose of rat PDGF-DD is typically less than 0.039 ng/ml. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined the mean O.D value of 20 replicates of the zero standard added by their three standard deviations.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess.
Inter-assay Precision	Inter-assay Precision (Precision between assays): CV%<10%. Three samples of known concentration were tested in twenty assays to assess.
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PDGF-DD elisa kit	Principle of the assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for PDGF-DD has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any PDGF-DD present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PDGF-DD is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PDGF-DD bound in the initial step. The color development is stopped and the intensity of the color is measured.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	75044560
UniProt Accession #	Q6V9H4
Molecular Weight	34,616 Da
NCBI Official Full Name	Platelet-derived growth factor D
UniProt Protein Name	Platelet-derived growth factor D
UniProt Gene Name	PDGFD
UniProt Synonym Gene Names	IEGF; PDGF-D; PDGFD latent form; PDGFD receptor-binding form
UniProt Entry Name	PDGFD_RABIT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/pdgf-dd-rat-elisa-kits/platelet-derived-growth-factor-dd/18173>