

Bovine Plasminogen Activator, Urokinase (uPA) ELISA Kit | uPA elisa kit

Cattle Plasminogen Activator, Urokinase (uPA) ELISA Kit

CATALOG #
AAA183006

REACTIVITY
Bovine


PRODUCT SYNONYMS

Plasminogen Activator; Urokinase (uPA); Cattle Plasminogen Activator; Urokinase (uPA) ELISA Kit; PLAU; ATF; URK; UK; UP-A; Abbokinase; Urokinase-Type Plasminogen Activator; uPA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Bovine
Specificity	This assay has high sensitivity and excellent specificity for detection of uPA. No significant cross-reactivity or interference between uPA and analogues was observed.
Sequence Length	431
Assay Type	Quantitative Sandwich
Samples	Bovine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.114ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level uPA were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level uPA were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for uPA elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to uPA. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to uPA. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain uPA, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of uPA in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of uPA in bovine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	15488889
NCBI GeneID	5328
UniProt Accession #	P00749; Q15844; Q16618; Q53XS3; Q5SWW9; Q969W6; B4DPZ2
Molecular Weight	46,908 Da
NCBI Official Full Name	Plasminogen activator, urokinase
NCBI Official Synonym Full Names	plasminogen activator, urokinase
NCBI Official Symbol	PLAU
NCBI Official Synonym Symbols	ATF; QPD; UPA; URK; u-PA; BDPLT5
NCBI Protein Information	urokinase-type plasminogen activator
UniProt Protein Name	Urokinase-type plasminogen activator
UniProt Gene Name	PLAU
UniProt Synonym Gene Names	U-plasminogen activator; uPA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/bovine-ovine-elisa-kits/plasminogen-activator-urokinase-upa/183006>