

Mouse Vascular Cell Adhesion Molecule 1 (VCAM1) ELISA Kit | VCAM1 elisa kit

Mouse Vascular Cell Adhesion Molecule 1 (VCAM1) ELISA Kit

CATALOG #
AAA183539

REACTIVITY
Mouse


PRODUCT SYNONYMS

Vascular Cell Adhesion Molecule 1 (VCAM1); Mouse Vascular Cell Adhesion Molecule 1 (VCAM1) ELISA Kit; CD106; INCAM-100; L1CAM; VCAM1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of VCAM1. No significant cross-reactivity or interference between VCAM1 and analogues was observed.
Sequence Length	647
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.39-25ng/mL
Sensitivity	< 0.14ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level VCAM1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level VCAM1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for VCAM1 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of VCAM1 in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to VCAM1. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to VCAM1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain VCAM1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of VCAM1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	179886
NCBI GeneID	7412
UniProt Accession #	P19320; Q6NUP8; A8K6R7; B4DKS4; E9PDD1
Molecular Weight	74,346 Da
NCBI Official Full Name	vascular cell adhesion molecule 1
NCBI Official Synonym Full Names	vascular cell adhesion molecule 1
NCBI Official Symbol	VCAM1
NCBI Official Synonym Symbols	CD106; INCAM-100
NCBI Protein Information	vascular cell adhesion protein 1
UniProt Protein Name	Vascular cell adhesion protein 1
UniProt Gene Name	VCAM1
UniProt Synonym Gene Names	V-CAM 1; VCAM-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/vascular-cell-adhesion-molecule-1-vcam1/183539>