

Mouse Thymus Activation Regulated Chemokine (TARC) ELISA Kit | TARC elisa kit

Mouse Thymus Activation Regulated Chemokine (TARC) ELISA Kit

CATALOG #
AAA184838

REACTIVITY
Mouse


PRODUCT SYNONYMS

Thymus Activation Regulated Chemokine (TARC); Mouse Thymus Activation Regulated Chemokine (TARC) ELISA Kit; CCL17; ABCD-2; SCYA17; TARC; Small Inducible Cytokine Subfamily A(Cys-Cys)Member 17; Small-inducible cytokine A17; TARC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Thymus Activation Regulated Chemokine (TARC). No significant cross-reactivity or interference between Thymus Activation Regulated Chemokine (TARC) and analogues was observed.
Sequence Length	93
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and Other Biological Fluids
Detection Range	31.25-2000pg/mL
Sensitivity	< 11.3pg/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Thymus Activation Regulated Chemokine (TARC) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Thymus Activation Regulated Chemokine (TARC) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TARC elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Thymus Activation Regulated Chemokine (TARC). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Thymus Activation Regulated Chemokine (TARC). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Thymus Activation Regulated Chemokine (TARC), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Thymus Activation Regulated Chemokine (TARC) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	11127960
NCBI GeneID	117518
Molecular Weight	10,480 Da
NCBI Official Full Name	thymus and activation-regulated chemokine
NCBI Official Synonym Full Names	C-C motif chemokine ligand 17
NCBI Official Symbol	Ccl17
NCBI Official Synonym Symbols	Tarc; Scya17
NCBI Protein Information	C-C motif chemokine 17
UniProt Protein Name	C-C motif chemokine
UniProt Gene Name	Ccl17

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.