

Cellular tumor antigen p53 (Tp53) Recombinant Protein | Tp53 recombinant protein

Recombinant Mouse Cellular tumor antigen p53 (Tp53)

CATALOG # AAA18501	HOST E Coli or Yeast or Baculovirus or Mammalian Cell
------------------------------	---


PRODUCT SYNONYMS

Cellular tumor antigen p53 (Tp53); Recombinant Mouse Cellular tumor antigen p53 (Tp53); Tumor suppressor p53; Tp53 recombinant protein

PRODUCT SPECIFICATIONS

Host	E Coli or Yeast or Baculovirus or Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	1-387aa; Full Length
Sequence	MEESQSDISL ELPLSQETFS GLWKLLPPED ILPSPHCMDD LLLPQDVVEF FEGPSEALRV SGAPAAQDPV TETPGPVAPA PATPWPLSSF VPSQKTYQGN YGFHLGFLQS GTAKSVMCTY SPPLNKLFCQ LAKTCPVQLW VSATPPAGSR VRAMAIYKKS QHMTEVVRRC PHERCSDGD GLAPPQHLIR VEGNLYPEYL EDRQTFRHSV VVPYEPPEAG SEYTTIHYKY MCNSSCMGGM NRRPILTIIT LEDSSGNLLG RDSFEVRVCA CPGRDRRTEE ENFRKKEVLC PELPPGSAKR ALPTCTSASP PQKKKPLDGE YFTLKIRGRK RFEMFRELNE ALELKDAHAT EESGDSRAHS SYLKTCKGQS TSRHKKTVMK KVGPDSD
Preparation and Storage	Store at -20 degree C, for extended storage, conserve at -20 degree C or -80 degree C.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for Tp53 recombinant protein	Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction ses to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis, but ses to have to effect on cell-cycle regulation. Regulates the circadian clock by repressing CLOCK-ARNTL/BMAL1-mediated transcriptional activation of PER2.
--	---

References

- Analysis of the gene coding for the murine cellular tumour antigen p53. Bienz B., Zakut-Houri R., Givol D., Oren M. *EMBO J.* 3:2179-2183(1984) A single gene and a pseudogene for the cellular tumour antigen p53. Zakut-Houri R., Oren M., Bienz B., Lavie V., Hazum S., Givol D. *Nature* 306:594-597(1983) Cloning and expression analysis of full length mouse cDNA sequences encoding the transformation associated protein p53. Jenkins J.R., Rudge K., Redmond S., Wade-Evans A. *Nucleic Acids Res.* 12:5609-5626(1984) Immunologically distinct p53 molecules generated by alternative splicing. Arai N., Nomura D., Yokota K., Wolf D., Brill E., Shohat O., Rotter V. *Mol. Cell. Biol.* 6:3232-3239(1986) Primary structure of DNA complementary to mRNA of murine oncoprotein p53. Chumakov P.M. *Bioorg. Khim.* 13:1691-1694(1987) Cell cycle in DNA-PKcs knock-out mice. Araki R., Fukumura R., Fujimori A., Tatsumi K., Abe M. DNA-dependent protein kinase is not required for the p53-dependent response to DNA damage. Jimenez G.S., Bryntesson F., Torres-Arzayus M.I., Priestley A., Beeche M., Saito S., Sakaguchi K., Appella E., Jeggo P.A., Taccioli G.E., Wahl G.M., Hubank M. *Nature* 400:81-83(1999) Characterization of DNA-PKcs null mutant SX9. Araki R., Fukumura R., Fujimori A., Tatsumi K., Abe M. p53 in 129-SVJ mice. Fujimori A., Abe M. Loss of heterozygosity and mutational alterations of the p53 gene in skin tumours of interspecific hybrid mice. Burns P.A., Kemp C.J., Gannon J.V., Lane D.P., Bremner R., Balmain A. *Oncogene* 6:2363-2369(1991) Detection of a transformation-related antigen in chemically induced sarcomas and other transformed cells of the mouse. DeLeo A.B., Jay G., Appella E., Dubois G.C., Law L.W., Old L.J. *Proc. Natl. Acad. Sci. U.S.A.* 76:2420-2424(1979) p53 is involved in the p120E4F-mediated growth arrest. Sandy P., Gostissa M., Fogal V., Cecco L.D., Szalay K., Rooney R.J., Schneider C., Del Sal G. *Oncogene* 19:188-199(2000) Characterization of cells and gene-targeted mice deficient for the p53-binding kinase homeodomain-interacting protein kinase 1 (HIPK1). Kondo S., Lu Y., Debbas M., Lin A.W., Sarosi I., Itie A., Wakeham A., Tuan J., Saris C., Elliott G., Ma W., Benchimol S., Lowe S.W., Mak T.W., Thukral S.K. *Proc. Natl. Acad. Sci. U.S.A.* 100:5431-5436(2003) Axin stimulates p53 functions by activation of HIPK2 kinase through multimeric complex formation. Rui Y., Xu Z., Lin S., Li Q., Rui H., Luo W., Zhou H.-M., Cheung P.-Y., Wu Z., Ye Z., Li P., Han J., Lin S.-C. *EMBO J.* 23:4583-4594(2004) Mapping of phosphomonoester and apparent phosphodiester bonds of the oncogene product p53 from simian virus 40-transformed 3T3 cells. Samad A., Anderson C.W., Carroll R.B. *Proc. Natl. Acad. Sci. U.S.A.* 83:897-901(1986) The p53 tumour suppressor protein is phosphorylated at serine 389 by casein kinase II. Meek D.W., Simon S., Kikkawa U., Eckhart W. *EMBO J.* 9:3253-3260(1990) p53 is covalently linked to 5.8S rRNA. Fontoura B.M., Sorokina E.A., David E., Carroll R.B. *Mol. Cell. Biol.* 12:5145-5151(1992) Negative control of p53 by Sir2alpha promotes cell survival under stress. Luo J., Nikolaev A.Y., Imai S., Chen D., Su F., Shiloh A., Guarente L., Gu W. *Cell* 107:137-148(2001) Identification and characterization of murine mHAUSP encoding a deubiquitinating enzyme that regulates the status of p53 ubiquitination. Lim S.-K., Shin J.-M., Kim Y.-S., Baek K.-H. *Int. J. Oncol.* 24:357-364(2004) Differential effect of ik3-1/cables on p53- and p73-induced cell death. Tsuji K., Mizumoto K., Yamochi T., Nishimoto I., Matsuoka M.J. *Biol. Chem.* 277:2951-2957(2002) Direct interaction with and activation of p53 by SMAR1 retards cell-cycle progression at G2/M phase and delays tumor growth in mice. Kaul R., Mukherjee S., Ahmed F., Bhat M.K., Chhipa R., Galande S., Chattopadhyay S. *Int. J. Cancer* 103:606-615(2003) PML regulates p53 stability by sequestering Mdm2 to the nucleolus. Bernardi R., Scaglioni P.P., Bergmann S., Horn H.F., Vousden K.H., Pandolfi P.P. *Nat. Cell Biol.* 6:665-672(2004) PRAK is essential for ras-induced senescence and tumor suppression. Sun P., Yoshizuka N., New L., Moser B.A., Li Y., Liao R., Xie C., Chen J., Deng Q., Yamout M., Dong M.Q., Frangou C.G., Yates J.R. III, Wright P.E., Han J. *Cell* 128:295-308(2007) Hzf Determines cell survival upon genotoxic stress by modulating p53 transactivation. Das S., Raj L., Zhao B., Kimura Y., Bernstein A., Aaronson S.A., Lee S.W. *Cell* 130:624-637(2007) Novel homeodomain-interacting protein kinase family member, HIPK4, phosphorylates human p53 at serine 9. Arai S., Matsushita A., Du K., Yagi K., Okazaki Y., Kurokawa R. *FEBS Lett.* 581:5649-5657(2007) CHD8 suppresses p53-mediated apoptosis through histone H1 recruitment during early embryogenesis. Nishiyama M., Oshikawa K., Tsukada Y.-I., Nakagawa T., Iemura S., Natsume T., Fan Y., Kikuchi A., Skoultschi A.I., Nakayama K.I. *Nat. Cell Biol.* 11:172-182(2009) Trim24 targets endogenous p53 for degradation. Allton K., Jain A.K., Herz H.M., Tsai W.W., Jung S.Y., Qin J., Bergmann A., Johnson R.L., Barton M.C. *Proc. Natl. Acad. Sci. U.S.A.* 106:11612-11616(2009) A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. Huarte M., Guttman M., Feldser D., Garber M., Koziol M.J., Kenzelmann-Broz D., Khalil A.M., Zuk O., Amit I., Rabani M., Attardi L.D., Regev A., Lander E.S., Jacks T., Rinn J.L. *Cell* 142:409-419(2010) p53 opens the mitochondrial permeability transition pore to trigger necrosis. Vaseva A.V., Marchenko N.D., Ji K., Tsirka S.E., Holzmans S., Moll U.M. *Cell* 149:1536-1548(2012) SIRT5-mediated lysine desuccinylation impacts diverse metabolic pathways. Park J., Chen Y., Tishkoff D.X., Peng C., Tan M., Dai L., Xie Z., Zhang Y., Zwaans B.M., Skinner M.E., Lombard D.B., Zhao Y. *Mol. Cell* 50:919-930(2013) p53 regulates Period2 expression and the circadian clock. Miki T., Matsumoto T., Zhao Z., Lee C.C. *Nat. Commun.* 4:2444-2444(2013) Crystal structure of the mouse p53 core DNA-binding domain at 2.7 Å resolution. Zhao K., Chai X., Johnston K., Clements A., Marmorstein R.J. *Biol. Chem.* 276:12120-12127(2001) High-resolution structure of the p53 core domain implications for binding small-molecule stabilizing compounds. Ho W.C., Luo C., Zhao K., Chai X., Fitzgerald M.X., Marmorstein R. *Acta Crystallogr. D* 62:1484-1493(2006) Structure of the p53 core domain dimer bound to DNA. Ho W.C., Fitzgerald M.X., Marmorstein R.J. *Biol. Chem.* 281:20494-20502(2006) Crystal structure of the mouse p53 core domain in zinc-free state. Kwon E., Kim D.Y., Suh S.W., Kim K.K. *Proteins* 70:280-283(2008) Crystal structure of a p53 core tetramer bound to DNA. Malecka K.A., Ho W.C., Marmorstein R. *Oncogene* 28:325-333(2009)

NCBI AND UNIPROT INFORMATION

NCBI GI #	187960040
NCBI GeneID	22059
NCBI Accession #	NP_001120705.1
NCBI GenBank Nucleotide #	NM_001127233.1
UniProt Accession #	P02340; Q9QUP3
Molecular Weight	59.2 kDa
NCBI Official Full Name	cellular tumor antigen p53 isoform b
NCBI Official Synonym Full Names	transformation related protein 53
NCBI Official Symbol	Trp53
NCBI Official Synonym Symbols	bbl; bfy; bhy; p44; p53; Tp53
NCBI Protein Information	cellular tumor antigen p53
UniProt Protein Name	Cellular tumor antigen p53
UniProt Gene Name	Tp53
UniProt Synonym Gene Names	P53; Trp53
UniProt Entry Name	P53_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/tp53-recombinant-protein-e-coli/18501>