

Human Nucleolin (NCL) ELISA Kit | NCL elisa kit

Human Nucleolin (NCL) ELISA Kit

CATALOG #
AAA185621

REACTIVITY
Human


PRODUCT SYNONYMS

Nucleolin (NCL); Human Nucleolin (NCL) ELISA Kit; C23; Protein C23; NCL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Nucleolin (NCL). No significant cross-reactivity or interference between Nucleolin (NCL) and analogues was observed.
Sequence Length	710
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.055ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Nucleolin (NCL) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Nucleolin (NCL) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NCL elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Nucleolin (NCL). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Nucleolin (NCL). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Nucleolin (NCL), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Nucleolin (NCL) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	55956788
NCBI GeneID	4691
NCBI Accession #	NP_005372.2
NCBI GenBank Nucleotide #	NM_005381.2
UniProt Accession #	P19338; Q53SK1; Q8NB06; Q9UCF0; Q9UDG1
Molecular Weight	76,614 Da
NCBI Official Full Name	nucleolin
NCBI Official Synonym Full Names	nucleolin
NCBI Official Symbol	NCL
NCBI Official Synonym Symbols	C23; Nsr1
NCBI Protein Information	nucleolin
UniProt Protein Name	Nucleolin
UniProt Gene Name	NCL

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.