

Human Platelet Activating Factor Receptor (PAFR) ELISA Kit | PAFR elisa kit

Human Platelet Activating Factor Receptor (PAFR) ELISA Kit

CATALOG #
AAA185912

REACTIVITY
Human


PRODUCT SYNONYMS

Platelet Activating Factor Receptor (PAFR); Human Platelet Activating Factor Receptor (PAFR) ELISA Kit; PTAFR; PAFR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Platelet Activating Factor Receptor (PAFR). No significant cross-reactivity or interference between Platelet Activating Factor Receptor (PAFR) and analogues was observed.
Sequence Length	342
Assay Type	Double-antibody Sandwich
Samples	Tissue Homogenates, Cell Lysates and Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.055ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Platelet Activating Factor Receptor (PAFR) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Platelet Activating Factor Receptor (PAFR) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PAFR elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Platelet Activating Factor Receptor (PAFR). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Platelet Activating Factor Receptor (PAFR). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Platelet Activating Factor Receptor (PAFR), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Platelet Activating Factor Receptor (PAFR) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506241
NCBI GeneID	5724
NCBI Accession #	NP_000943.1
NCBI GenBank Nucleotide #	NM_000952.4
UniProt Accession #	P25105; A3KMC8; A8K2H5
Molecular Weight	39,203 Da
NCBI Official Full Name	platelet-activating factor receptor
NCBI Official Synonym Full Names	platelet activating factor receptor
NCBI Official Symbol	PTAFR
NCBI Official Synonym Symbols	PAFR
NCBI Protein Information	platelet-activating factor receptor
UniProt Protein Name	Platelet-activating factor receptor
UniProt Gene Name	PTAFR
UniProt Synonym Gene Names	PAFR; PAF-R; PAFr

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/platelet-activating-factor-receptor-pafr/185912>