

Rat Interleukin 20 (IL20) CLIA Kit | IL20 clia kit

Rat Interleukin 20 (IL20) CLIA Kit

CATALOG # AAA189072	REACTIVITY Rat	APPLICATIONS CLIA (Chemiluminescence Immunoassay)
-------------------------------	--------------------------	---


PRODUCT SYNONYMS

Interleukin 20 (IL20); Rat Interleukin 20 (IL20) CLIA Kit; Interleukin 20; IL10D; IL10-D; ZCYTO10; Cytokine Zcyto10; IL20 clia kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Interleukin 20 (IL20). No significant cross-reactivity or interference between Interleukin 20 (IL20) and analogues was observed.
Sequence Length	176
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	1.37-1000pg/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 0.54pg/mL
Applicable Applications for IL20 clia kit	CLIA (Chemiluminescence Immunoassay)
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Interleukin 20 (IL20) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Interleukin 20 (IL20) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The microplate provided in this kit has been pre-coated with an antibody specific to Interleukin 20 (IL20). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Interleukin 20 (IL20). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Interleukin 20 (IL20) level in the sample or standard.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	219842195
NCBI GeneID	681112
NCBI Accession #	NP_001137353.1
NCBI GenBank Nucleotide #	NP_001137353.1
UniProt Accession #	D4A2T8
NCBI Official Full Name	interleukin-20
NCBI Official Synonym Full Names	interleukin 20
NCBI Official Symbol	IL20
NCBI Protein Information	interleukin-20
UniProt Protein Name	Protein IL20
UniProt Gene Name	IL20

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/rat-clia-kits/interleukin-20-il20/189072>