

Rabbit anti-Human SOCS1 Polyclonal Antibody | anti-SOCS1 antibody

Anti-SOCS1 Antibody Picoband


CATALOG # AAA19396	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---

PRODUCT SYNONYMS

SOCS1; Polyclonal Antibody; Anti-SOCS1 Antibody Picoband; anti-SOCS1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-SOCS1 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Application Notes	WB: 0.25-0.5 ug/ml, Human IHC-P: 2-5 ug/ml, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC(P).
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human SOCS1 (FLVRDSRQRNCFALSVKMA).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 6. IHC analysis of SOCS1 using anti-SOCS1 antibody (AAA19396).SOCS1 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOCS1 Antibody (AAA19396) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 5. IHC analysis of SOCS1 using anti-SOCS1 antibody (AAA19396).SOCS1 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOCS1 Antibody (AAA19396) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 4. IHC analysis of SOCS1 using anti-SOCS1 antibody (AAA19396).SOCS1 was detected in a paraffin-embedded section of human esophageal squamous carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOCS1 Antibody (AAA19396) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of SOCS1 using anti-SOCS1 antibody (AAA19396).SOCS1 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOCS1 Antibody (AAA19396) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of SOCS1 using anti-SOCS1 antibody (AAA19396).SOCS1 was detected in a paraffin-embedded section of human cervical cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOCS1 Antibody (AAA19396) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of SOCS1 using anti-SOCS1 antibody (AAA19396).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human K562 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SOCS1 antigen affinity purified polyclonal antibody (#AAA19396) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for SOCS1 at approximately 27 kDa. The expected band size for SOCS1 is at 24 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-SOCS1 antibody	Suppressor of cytokine signaling 1(SOCS1), also known as Jak-binding protein(JAB), is a negative regulator to a subset of protein-tyrosine kinases. This 211-amino acid protein has a molecular mass of about 23.5 kD. The TEC-binding region of SOCS1 resides in the N terminus. SOCS1 associates with Tec and suppresses its kinase activity. The SOCS1 gene is intronless and is mapped to on chromosome 16p13.13. SOCS1 regulates the JAK/STAT signal-transduction pathway. Moreover, it contributes to protection against hepatic injury and fibrosis, and may also protect against hepatocarcinogenesis.
Product Categories/Family for anti-SOCS1 antibody	Primary Antibodies , Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507233
NCBI GeneID	8651
NCBI Accession #	NP_003736.1
NCBI GenBank Nucleotide #	NM_003745.1
UniProt Accession #	O15524 ; O15097 ; Q9NSA7
Molecular Weight	211
NCBI Official Full Name	suppressor of cytokine signaling 1
NCBI Official Synonym Full Names	suppressor of cytokine signaling 1
NCBI Official Symbol	SOCS1
NCBI Official Synonym Symbols	JAB; CIS1; SSI1; TIP3; CISH1; SSI-1; SOCS-1
NCBI Protein Information	suppressor of cytokine signaling 1; TIP-3; JAK binding protein; JAK-binding protein; Tec-interacting protein 3; STAT induced SH3 protein 1; STAT-induced STAT inhibitor 1; cytokine-inducible SH2 protein 1
UniProt Protein Name	Suppressor of cytokine signaling 1

UniProt Gene Name	SOCS1
UniProt Synonym Gene Names	SSI1; TIP3; SOCS-1; JAB; SSI-1; TIP-3
UniProt Entry Name	SOCS1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/socs1-antibody-western-blot-immunohistochemistry-elisa/19396>