

Mouse PDIA6 Monoclonal Antibody | anti-PDIA6 antibody

Anti-PDIA6 Antibody Picoband (monoclonal, 3H5E7)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19710	Mouse	Human, Monkey	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

PDIA6; Monoclonal Antibody; Anti-PDIA6 Antibody Picoband (monoclonal; 3H5E7); anti-PDIA6 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Monkey
Clonality	Monoclonal
Isotype	Mouse IgG2b
Clone Number	3H5E7
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-PDIA6 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Application Notes	WB: 0.25-0.5 ug/ml, Human, Monkey IHC-P: 2-5 ug/ml, Human ICC/IF: 5 ug/ml, Human FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Mouse IgG for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit for IHC(P) and ICC.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human PDIA6 recombinant protein (Position: L20-L440).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM (Flow Cytometry)

Figure 11. Flow Cytometry analysis of SiHa cells using anti-PDIA6 antibody (AAA19710).Overlay histogram showing SiHa cells stained with AAA19710 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-PDIA6 Antibody (AAA19710, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 9. IHC analysis of PDIA6 using anti-PDIA6 antibody (AAA19710).PDIA6 was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-PDIA6 Antibody (AAA19710) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.




IF (Immunofluorescence)

Figure 10. IF analysis of PDIA6 using anti-PDIA6 antibody (AAA19710).PDIA6 was detected in an immunocytochemical section of HepG2 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL mouse anti-PDIA6 Antibody (AAA19710) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IHC (Immunohistochemistry)

Figure 8. IHC analysis of PDIA6 using anti-PDIA6 antibody (AAA19710).PDIA6 was detected in a paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-PDIA6 Antibody (AAA19710) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 7. IHC analysis of PDIA6 using anti-PDIA6 antibody (AAA19710).PDIA6 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-PDIA6 Antibody (AAA19710) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 6. IHC analysis of PDIA6 using anti-PDIA6 antibody (AAA19710).PDIA6 was detected in a paraffin-embedded section of human laryngeal squamous cell carcinomas tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-PDIA6 Antibody (AAA19710) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-PDIA6 antibody	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, two catalytically active thioredoxin (TRX) domains, a TRX-like domain, and a C-terminal ER-retention sequence. This protein inhibits the aggregation of misfolded proteins and exhibits both isomerase and chaperone activity. Alternative splicing results in multiple transcript variants encoding different isoforms.
Product Categories/Family for anti-PDIA6 antibody	Primary Antibodies , Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	2501205
NCBI GeneID	10130
UniProt Accession #	Q15084 ; Q53RC7 ; Q6ZSH5 ; Q99778 ; B3KY95 ; B5MCQ5 ; B7Z254 ; B7Z4M8 ; F8WA83
Molecular Weight	440
NCBI Official Full Name	Protein disulfide-isomerase A6
NCBI Official Synonym Full Names	protein disulfide isomerase family A, member 6
NCBI Official Symbol	PDIA6
NCBI Official Synonym Symbols	P5; ERP5; TXNDC7
NCBI Protein Information	protein disulfide-isomerase A6; ER protein 5; protein disulfide isomerase P5; endoplasmic reticulum protein 5; thioredoxin domain-containing protein 7; protein disulfide isomerase-associated 6; protein disulfide isomerase-related protein; thioredoxin doma
UniProt Protein Name	Protein disulfide-isomerase A6
UniProt Gene Name	PDIA6
UniProt Synonym Gene Names	ERP5; P5; TXNDC7; ER protein 5; ERp5
UniProt Entry Name	PDIA6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

