

Rabbit LTA4H Polyclonal Antibody | anti-LTA4H antibody

Anti-LTA4H Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19774	Rabbit	Human, Mouse, Rat	ELISA, Immunofluorescence, Immunohistochemistry, Western Blot, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

LTA4H; Polyclonal Antibody; Anti-LTA4H Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-LTA4H antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-LTA4H antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), FCM/FACS (Flow Cytometry)
Application Notes	WB: 0.1-0.25ug/ml, Human, Mouse, Rat IHC: 1-2ug/ml, Human IF: 5ug/ml, Human FC/FACS (Fixed): 1-3ug/1x10 ⁶ cells, Human ELISA: 0.1-0.5ug/ml
Immunogen	E coli-derived human IKZF3 recombinant protein (Position: M1-H456). Human IKZF3 shares 85.5% amino acid (aa) sequence identity with mouse IKZF3.
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



FCM (Flow Cytometry)

Figure 7. Flow Cytometry analysis of RT4 cells using anti-LTA4H antibody (AAA19774). Overlay histogram showing RT4 cells stained with AAA19774 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-LTA4H Antibody (AAA19774, 1ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 6. IF analysis of LTA4H using anti-LTA4H antibody (AAA19774). LTA4H was detected in a paraffin-embedded section of human intestinal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5ug/mL rabbit anti-LTA4H Antibody (AAA19774) overnight at 4 degree C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

NCBI GeneID	4048
UniProt Accession #	P09960; Q6IAT6; Q9UCT7; B4DNQ9; F8VV40
Molecular Weight	611
NCBI Official Full Name	Leukotriene A-4 hydrolase
NCBI Official Synonym Full Names	leukotriene A4 hydrolase
NCBI Official Symbol	LTA4H
NCBI Protein Information	leukotriene A-4 hydrolase; LTA-4 hydrolase
UniProt Protein Name	Leukotriene A-4 hydrolase
UniProt Gene Name	LTA4H
UniProt Synonym Gene Names	LTA4; LTA-4 hydrolase
UniProt Entry Name	LKHA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/lta4h-antibody-immunofluorescence-immunohistochemistry-western-blot-flow-cytometry-elisa/19774>