

Rabbit anti-Human MAPK8 Polyclonal Antibody | anti-MAPK8 antibody

MAPK8 Antibody - N-terminal region

CATALOG # AAA201559	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot
-------------------------------	-----------------------	----------------------------	-------------------------------------


PRODUCT SYNONYMS

MAPK8; Polyclonal Antibody; MAPK8 Antibody - N-terminal region; anti-MAPK8 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Affinity purified
Form/Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Sequence	Synthetic peptide with the following region: MS RSKRDNNFYS VEIGDSTFTV LKRYQNLKPI GSGAQGIVCA AYDAILER
Sequence Length	427
Applicable Applications for anti-MAPK8 antibody	WB (Western Blot)
Immunogen	The immunogen is a synthetic peptide directed towards the N terminal region of human MAPK8
Preparation and Storage	For short term use, store at 2-8 degree C up to 1 week. For long term storage, store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

Host: Rabbit
Target Name: MAPK8
Sample Tissue: Human DLD1 Whole Cell
Antibody Dilution: 1.0ug/ml


IHC (Immunohistochemistry)

Rabbit Anti-MAPK8 Antibody
Formalin Fixed Paraffin Embedded Tissue: Human Thyroid Gland Tissue
Observed Staining: Cytoplasm
Primary Antibody Concentration: 1:100
Other Working Concentrations: N/A
Secondary Antibody: Donkey anti-Rabbit-Cy3
Secondary Antibody Concentration: 1:200
Magnification: 20X
Exposure Time: 0.5 - 2.0 sec

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK8 antibody	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, apoptosis and differentiation. Five alternatively spliced transcript variants encoding distinct isoforms have been reported.
Product Categories/Family for anti-MAPK8 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	513788279
NCBI GeneID	5599
NCBI Accession #	NP_001265476.1
NCBI GenBank Nucleotide #	NM_001278547.1
UniProt Accession #	P45983
Molecular Weight	48 kDa
NCBI Official Full Name	mitogen-activated protein kinase 8 isoform beta2
NCBI Official Synonym Full Names	mitogen-activated protein kinase 8
NCBI Official Symbol	MAPK8
NCBI Official Synonym Symbols	JNK; JNK1; PRKM8; SAPK1; JNK-46; JNK1A2; SAPK1c; JNK21B1/2
NCBI Protein Information	mitogen-activated protein kinase 8
UniProt Protein Name	Mitogen-activated protein kinase 8
UniProt Gene Name	MAPK8
UniProt Synonym Gene Names	JNK1; PRKM8; SAPK1; SAPK1C; MAP kinase 8; MAPK 8; SAPK1c
UniProt Entry Name	MK08_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.