

Human Neprilysin (NEP) ELISA Kit | NEP elisa kit

Human Neprilysin (NEP) ELISA Kit

CATALOG #
AAA20416

REACTIVITY
Human


PRODUCT SYNONYMS

Neprilysin (NEP); Human Neprilysin (NEP) ELISA Kit; CD10; MME; CALLA; SFE; Atriopeptidase; Enkephalinase; Skin fibroblast elastase; Common Acute Lymphoblastic Leukemia Antigen; Membrane Metallo-Endopeptidase; NEP elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of neprilysin. No significant cross-reactivity or interference between neprilysin and analogues was observed.
Sequence Length	750
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, cerebrospinal fluid and other biological fluids
Detection Range	31.2-2,000pg/mL
Sensitivity	< 12.5pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level neprilysin were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level neprilysin were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NEP elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to neprilysin. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to neprilysin. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain neprilysin, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of neprilysin in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of neprilysin in human serum, plasma, cerebrospinal fluid and other biological fluids.
Product Categories/Family for NEP elisa kit	Enzyme & Kinase; Neuroscience

NCBI AND UNIPROT INFORMATION

NCBI GI #	386782299
NCBI GeneID	707667
NCBI Accession #	NP_001247743.1
NCBI GenBank Nucleotide #	NM_001260814.1
NCBI Official Full Name	neprilysin
NCBI Official Symbol	MME
NCBI Official Synonym Symbols	neprilysin
NCBI Protein Information	neprilysin

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/neprilysin-nep/20416>