

Mouse Cluster Of Differentiation 40 Ligand (CD40L) ELISA Kit | CD40L elisa kit

Mouse Cluster Of Differentiation 40 Ligand (CD40L) ELISA Kit

CATALOG #
AAA20747

REACTIVITY
Mouse


PRODUCT SYNONYMS

Cluster Of Differentiation 40 Ligand (CD40L); Mouse Cluster Of Differentiation 40 Ligand (CD40L) ELISA Kit; CD154; TRAP; HIGM1; IGM; IMD3; TBAM; T-BAM; TNFSF5; Gp39; TNF Superfamily Member 5; Hyper-IgM Syndrome; TNF-Related Activation Protein; T-Cell B-Cell Activating Molecule; CD40L elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CD40L. No significant cross-reactivity or interference between CD40L and analogues was observed.
Sequence Length	261
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.062ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CD40L were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CD40L were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD40L elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to CD40L. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to CD40L. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CD40L, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of CD40L in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CD40L in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Product Categories/Family for CD40L elisa kit	Signal transduction; CD & Adhesion molecule; Apoptosis; Tumor immunity; Infection immunity; Immune molecule

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557433
NCBI GeneID	959
NCBI Accession #	NP_000065.1
NCBI GenBank Nucleotide #	NM_000074.2
UniProt Accession #	P29965
Molecular Weight	29,274 Da
NCBI Official Full Name	CD40 ligand
NCBI Official Synonym Full Names	CD40 ligand
NCBI Official Symbol	CD40LG
NCBI Official Synonym Symbols	IGM; IMD3; TRAP; gp39; CD154; CD40L; HIGM1; T-BAM; TNFSF5; hCD40L
NCBI Protein Information	CD40 ligand
UniProt Protein Name	CD40 ligand
UniProt Gene Name	CD40LG
UniProt Synonym Gene Names	CD40L; TNFSF5; TRAP; CD40-L; TRAP
UniProt Entry Name	CD40L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.