

Rabbit anti-Human NRIP Polyclonal Antibody | anti-NRIP antibody

Rabbit anti-NRIP Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA211874	Rabbit	Human	Immunohistochemistry, Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

NRIP; Polyclonal Antibody; Rabbit anti-NRIP Antibody; Affinity Purified; IQ motif and WD repeat-containing protein 1; nuclear receptor interaction protein; MSTP055; IQ motif and WD repeats 1; PC326; DDB1- and CUL4-associated factor 6; ARCAP; androgen receptor complex-associated protein; 1200006M05Rik; IQWD1; DDB1 and CUL4 associated factor 6; anti-NRIP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	200ug/ml (varies by lot)
Applicable Applications for anti-NRIP antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Immunogen	Between 600 and 650
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human NRIP by western blot and immunoprecipitation. Samples: Whole cell lysate (5, 15 and 50 ug for WB; 1 mg for IP, 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti-NRIP antibody AAA211874 used for WB at 0.04 ug/ml (A) and 1 ug/ml (B) and used for IP at 10 ug/mg lysate. NRIP was also immunoprecipitated by rabbit anti-NRIP antibody which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 10 seconds (B).



IHC (Immunohistochemistry)

Detection of human NRIP by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Affinity purified rabbit anti-NRIP (Cat. No. AAA211874 Lot1) used at a dilution of 1:200 (1ug/ml). Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-NRIP antibody	Nuclear receptor interaction protein (NRIP) was isolated as a ligand-dependent coactivator of the androgen receptor (AR) and glucocorticoid receptor (GR). NRIP was found to enhance the promoter activity of the AR-driven PSA promoter. NRIP is a target gene for AR and Sp1, and complex formation between AR and NRIP protects AR protein from proteasome degradation.
Product Categories/Family for anti-NRIP antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	63252908
NCBI GeneID	55827
NCBI Accession #	NP_060912.2
NCBI GenBank Nucleotide #	NM_018442.3
UniProt Accession #	Q58WW2; Q7L8I0; Q8IXH3; Q8TB19; A2A295; B4DNB8
Molecular Weight	103,070 Da
NCBI Official Full Name	DDB1- and CUL4-associated factor 6 isoform a
NCBI Official Synonym Full Names	DDB1 and CUL4 associated factor 6
NCBI Official Symbol	DCAF6
NCBI Official Synonym Symbols	NRIP; ARCAP; IQWD1; PC326; MSTP055; 1200006M05Rik
NCBI Protein Information	DDB1- and CUL4-associated factor 6; IQ motif and WD repeats 1; nuclear receptor interaction protein; IQ motif and WD repeat-containing protein 1; androgen receptor complex-associated protein
UniProt Protein Name	DDB1- and CUL4-associated factor 6
UniProt Gene Name	DCAF6
UniProt Synonym Gene Names	IQWD1; ARCAP; NRIP
UniProt Entry Name	DCAF6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/nrip-antibody-immunohistochemistry-immunoprecipitation-western-blot-elisa/211874>