

Rabbit anti-Human PRRC2C Polyclonal Antibody | anti-PRRC2C antibody

Rabbit anti-PRRC2C Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA212150	Rabbit	Human	Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

PRRC2C; Polyclonal Antibody; Rabbit anti-PRRC2C Antibody; Affinity Purified; proline-rich and coiled-coil-containing protein 2C; HBV X-transactivated gene 2 protein; XTP2; protein BAT2-like 2; HLA-B-associated transcript 2-like 2; HBxAg transactivated protein 2; BAT2L2; BAT2-iso; BAT2 domain containing 1; BAT2D1; protein PRRC2C; BAT2 domain-containing protein 1; HBV XAg-transactivated protein 2; Protein PRRC2C; proline rich coiled-coil 2C; anti-PRRC2C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-PRRC2C antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	Between 2250 and 2300
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human PRRC2C by western blot. Samples: Whole cell lysate (5, 15 and 50 ug) from HeLa cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-PRRC2C antibody AAA212150 (lot AAA212150-2) used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.



IP (Immunoprecipitation)

Detection of human PRRC2C by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-PRRC2C antibody AAA212150 (lot AAA212150-2) used for IP at 6 ug per reaction. PRRC2C was also immunoprecipitated by rabbit anti-PRRC2C antibody For blotting immunoprecipitated PRRC2C, AAA212150 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.

ADDITIONAL INFORMATION

Related Product Information for anti-PRRC2C antibody	Proline-rich coiled-coil 2C (PRRC2C) is proposed to be involved in protein C-terminus binding and contains a conserved BAT2 N-terminus domain.
Product Categories/Family for anti-PRRC2C antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	115298682
NCBI GeneID	23215
NCBI Accession #	NP_055987.2
NCBI GenBank Nucleotide #	NM_015172.3
UniProt Accession #	Q9Y520; Q05DM8; Q49A39; Q6PD54; Q9H2N2; Q9HA05; Q9NSM8; Q9NXL3; Q9UF29; Q9UPQ6
Molecular Weight	316,911 Da
NCBI Official Full Name	protein PRRC2C
NCBI Official Synonym Full Names	proline-rich coiled-coil 2C
NCBI Official Symbol	PRRC2C
NCBI Official Synonym Symbols	XTP2; BAT2D1; BAT2L2; BAT2-iso
NCBI Protein Information	protein PRRC2C; protein BAT2-like 2; BAT2 domain containing 1; HBxAg transactivated protein 2; BAT2 domain-containing protein 1; HBV XAg-transactivated protein 2; HBV X-transactivated gene 2 protein; HLA-B associated transcript 2-like 2; HLA-B-associated
UniProt Protein Name	Protein PRRC2C
UniProt Gene Name	PRRC2C
UniProt Synonym Gene Names	BAT2D1; BAT2L2; KIAA1096; XTP2
UniProt Entry Name	PRC2C_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/prrc2c-antibody-immunoprecipitation-western-blot-elisa/212150>