

Rabbit anti-Human COMT Polyclonal Antibody | anti-COMT antibody

Rabbit anti-COMT Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA212245	Rabbit	Human	Immunohistochemistry, Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

COMT; Polyclonal Antibody; Rabbit anti-COMT Antibody; Affinity Purified; catechol O-methyltransferase; catechol-O-methyltransferase isoform; epididymis secretory sperm binding protein Li 98n; HEL-S-98n; testicular tissue protein Li 42; Catechol O-methyltransferase; catechol-O-methyltransferase; anti-COMT antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-COMT antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Immunogen	Between 221 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human COMT by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (15 and 50 ug for WB; 1 mg for IP, 20% of IP loaded), HEK293T (T; 50 ug) and Jurkat (J; 50 ug) cells. Antibodies: Affinity purified rabbit anti-COMT antibody AAA212245 used for WB at 0.1 ug/ml (A) and 1 ug/ml (B) and used for IP at 6 ug/mg lysate. COMT was also immunoprecipitated by rabbit anti-COMT antibody BL13028, which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 10 seconds (B).



IHC (Immunohistochemistry)

Detection of human COMT by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Affinity purified rabbit anti-COMT (Cat. No. AAA212245 Lot1) used at a dilution of 1:1,000 (1ug/ml). Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-COMT antibody	Catechol O-methyltransferase (COMT) catalyzes the O-methylation, and thereby the inactivation, of catecholamine neurotransmitters and catechol hormones. Also shortens the biological half-lives of certain neuroactive drugs, like L-DOPA, alpha-methyl DOPA and isoproterenol [taken from the Universal Protein Resource (UniProt) www.uniprot.org/uniprot/P21964].
Product Categories/Family for anti-COMT antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502969
NCBI GeneID	1312
NCBI Accession #	NP_000745.1
NCBI GenBank Nucleotide #	NM_000754.3
UniProt Accession #	P21964 ; Q6IB07 ; Q6ICE6 ; A8MPV9
Molecular Weight	30,037 Da
NCBI Official Full Name	catechol O-methyltransferase isoform MB-COMT
NCBI Official Synonym Full Names	catechol-O-methyltransferase
NCBI Official Symbol	COMT
NCBI Protein Information	catechol O-methyltransferase
UniProt Protein Name	Catechol O-methyltransferase
UniProt Gene Name	COMT
UniProt Entry Name	COMT_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/comt-antibody-immunohistochemistry-immunoprecipitation-western-blot-elisa/212245>