

Rabbit anti-Human TTC15 Polyclonal Antibody | anti-TTC15 antibody

Rabbit anti-TTC15 Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA212850	Rabbit	Human	Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

TTC15; Polyclonal Antibody; Rabbit anti-TTC15 Antibody; Affinity Purified; CGI-87; trafficking of membranes and mitosis; TTC-15; trafficking protein particle complex subunit 12; TPR repeat protein 15; tetratricopeptide repeat protein 15; PEBAS; tetratricopeptide repeat domain 15; Trafficking protein particle complex subunit 12; trafficking protein particle complex 12; anti-TTC15 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-TTC15 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	Between 75 and 125
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human TTC15 by western blot. Samples: Whole cell lysate (50 ug) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-TTC15 antibody AAA212850 (lot AAA212850-1) used for WB at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



IP (Immunoprecipitation)

Detection of human TTC15 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-TTC15 antibody AAA212850 (lot AAA212850-1) used for IP at 6 ug per reaction. TTC15 was also immunoprecipitated by rabbit anti-TTC15 antibody BL17979. For blotting immunoprecipitated TTC15, AAA212850 was used at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.

ADDITIONAL INFORMATION

Related Product Information for anti-TTC15 antibody	Tetratricopeptide repeat domain 15 (TTC15) may be involved in endoplasmic reticulum to Golgi apparatus trafficking at a very early stage [taken from the Universal Protein Resource (UniProt) www.uniprot.org/uniprot/Q8WVT3].
Product Categories/Family for anti-TTC15 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GeneID	51112
UniProt Accession #	Q8WVT3
UniProt Protein Name	Trafficking protein particle complex subunit 12
UniProt Gene Name	TRAPPC12
UniProt Synonym Gene Names	TTC15; TPR repeat protein 15; TTC-15
UniProt Entry Name	TPC12_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/ttc15-antibody-immunoprecipitation-western-blot-elisa/212850>