

Rabbit anti-Human ZFR Polyclonal Antibody | anti-ZFR antibody

Rabbit anti-ZFR Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA212985	Rabbit	Human	Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

ZFR; Polyclonal Antibody; Rabbit anti-ZFR Antibody; Affinity Purified; SPG71; ZFR1; hZFR; M-phase phosphoprotein homolog; zinc finger RNA-binding protein; Zinc finger RNA-binding protein; zinc finger RNA binding protein; anti-ZFR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-ZFR antibody	IP (Immunoprecipitation), WB (Western Blot)
Predicted Reactivity	Mouse, Rat, Zebrafish, Orangutan
Immunogen	Between 1 and 50
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human ZFR by western blot. Samples: Whole cell lysate (50 ug) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-ZFR antibody AAA212985 (lot AAA212985-1) used for WB at 0.4 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



IP (Immunoprecipitation)

Detection of human ZFR by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-ZFR antibody AAA212985 (lot AAA212985-1) used for IP at 6 ug per reaction. ZFR was also immunoprecipitated by rabbit anti-ZFR antibody BL18962. For blotting immunoprecipitated ZFR, AAA212985 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.

ADDITIONAL INFORMATION

Related Product Information for anti-ZFR antibody	Zinc finger RNA-binding protein (ZFR) is involved in postimplantation and gastrulation stages of developmen. Itis Involved in the nucleocytoplasmic shuttling of STAU2. Binds to DNA and RNA (By similarity) [taken from the Universal Protein Resource (UniProt) www.uniprot.org/uniprot/Q96KR1].
Product Categories/Family for anti-ZFR antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	34101286
NCBI GeneID	51663
NCBI Accession #	NP_057191.2
NCBI GenBank Nucleotide #	NM_016107.3
UniProt Accession #	Q96KR1; Q05C08; Q3B7X5; Q6P5A3; Q86UA0; Q9H6V4; Q9NTI1; Q9Y687; B2RNR5
Molecular Weight	117,012 Da
NCBI Official Full Name	zinc finger RNA-binding protein
NCBI Official Synonym Full Names	zinc finger RNA binding protein
NCBI Official Symbol	ZFR
NCBI Official Synonym Symbols	ZFR1; SPG71
NCBI Protein Information	zinc finger RNA-binding protein; hZFR; M-phase phosphoprotein homolog
UniProt Protein Name	Zinc finger RNA-binding protein
UniProt Gene Name	ZFR
UniProt Synonym Gene Names	hZFR
UniProt Entry Name	ZFR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/zfr-antibody-immunoprecipitation-western-blot-elisa/212985>