

Rabbit anti-Human, Mouse MED18 Polyclonal Antibody | anti-MED18 antibody

Rabbit anti-MED18 IHC Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA213852	Rabbit	Human, Mouse	Immunohistochemistry


PRODUCT SYNONYMS

MED18; Polyclonal Antibody; Rabbit anti-MED18 IHC Antibody; Affinity Purified; p28b; SRB5; mediator of RNA polymerase II transcription; subunit 18 homolog; mediator of RNA polymerase II transcription subunit 18; Mediator complex subunit 18; TRAP/mediator complex subunit p28b; Mediator of RNA polymerase II transcription subunit 18; mediator complex subunit 18; anti-MED18 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	250ug/ml (varies by lot)
Applicable Applications for anti-MED18 antibody	IHC (Immunohistochemistry)
Predicted Reactivity	Rat, Zebrafish, Bovine, Dog, Horse, Rabbit, Pig, Panda, Orangutan, Monkey, Gorilla, Chimpanzee, Crab-eating macaque, Duckbill platypus, White-tufted-ear marmoset, Tasmanian devil, Small-eared galago, Thirteen-lined ground squirrel, Gray short-tailed opossum, Zebra finch, Naked mole rat, Channel catfish, African elephant, Teleostei, Northern white-cheeked gibbon, Chinese hamster
Immunogen	Between 158 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



IHC (Immunohistochemistry)
Detection of human MED18 by immunohistochemistry. Sample: FFPE section of human prostate adenocarcinoma. Antibody: Affinity purified rabbit anti-MED18 (Cat. No. AAA213852) used at a dilution of 1:250. Detection: DAB



IHC (Immunohistochemistry)
Detection of mouse MED18 by immunohistochemistry. Sample: FFPE section of mouse squamous cell carcinoma. Antibody: Affinity purified rabbit anti-MED18 (Cat. No. AAA213852) used at a dilution of 1:250. Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-MED18 antibody	MED18 is a component of the multisubunit mediator complex that functions as a bridge between gene-specific transcription factors and the RNA Pol II/general transcription factor machinery. The mediator complex was first identified as a coactivator of the thyroid hormone receptor and has since been observed to function as a coactivator of a variety of transcription factors. MED18 forms a subcomplex with the MED8 and MED20 subunits and plays an important role in initiation complex formation.
Product Categories/Family for anti-MED18 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GeneID	54797
UniProt Accession #	Q9BUE0
UniProt Protein Name	Mediator of RNA polymerase II transcription subunit 18
UniProt Gene Name	MED18
UniProt Entry Name	MED18_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/med18-antibody-immunohistochemistry-elisa/213852>