

Rabbit anti-Human BCR Polyclonal Antibody | anti-BCR antibody

Rabbit anti-BCR IHC Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214006	Rabbit	Human	Immunocytochemistry, Immunocytochemistry

PRODUCT SYNONYMS

BCR; Polyclonal Antibody; Rabbit anti-BCR IHC Antibody; Affinity Purified; D22S11; PHL; BCR/FGFR1 chimera protein; FGFR1/BCR chimera protein; renal carcinoma antigen NY-REN-26; D22S662; breakpoint cluster region protein; BCR1; ALL; breakpoint cluster region; CML; Breakpoint cluster region protein; RhoGEF and GTPase activating protein; anti-BCR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	250ug/ml (varies by lot)
Applicable Applications for anti-BCR antibody	ICC (Immunocytochemistry), ICC (Immunocytochemistry)
Predicted Reactivity	Gorilla
Immunogen	Between 75 and 125
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



ICC (Immunocytochemistry)

Detection of human BCR by immunocytochemistry. Sample: FFPE section of human K-562 cells (contains the chromosomal translocation, t(9:22) that creates the BCR/ABL fusion gene). Antibody: Affinity purified rabbit anti-BCR (Cat. No. AAA214006) used at a dilution of 1:250. Detection: DAB



ICC (Immunocytochemistry)

Detection of human BCR by immunocytochemistry. Sample: FFPE section of human K-562 cells (contains the chromosomal translocation, t(9:22) that creates the BCR/ABL fusion gene). Antibody: Affinity purified rabbit anti-BCR (Cat. No. AAA214006) used at a dilution of 1:250. Detection: Red-fluorescent goat anti-rabbit IgG highly cross-adsorbed Antibody Hilyte Plus 555 used at a dilution of 1:100.

ADDITIONAL INFORMATION

Related Product Information for anti-BCR antibody	BCR (breakpoint cluster region) was identified as part of a fusion protein that results from a reciprocal translocation with the ABL gene between chromosomes 22 and 9. The BCR-ABL fusion is the cause of chronic myeloid leukemia (CML). The BCR protein contains a C2 domain, a DBL-homology domain, a PH domain, and a Rho-GAP domain and functions as a GTPase-activating protein for RAC1 and CDC42.
Product Categories/Family for anti-BCR antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GeneID	613
UniProt Accession #	P11274
UniProt Protein Name	Breakpoint cluster region protein
UniProt Gene Name	BCR
UniProt Synonym Gene Names	BCR1; D22S11
UniProt Entry Name	BCR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/bcr-antibody-immunocytochemistry-immunocytochemistry-elisa/214006>