

Rabbit anti-Human WDR20 Polyclonal Antibody | anti-WDR20 antibody

Rabbit anti-WDR20 IHC Antibody, Affinity Purified

CATALOG # AAA214101	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

WDR20; Polyclonal Antibody; Rabbit anti-WDR20 IHC Antibody; Affinity Purified; DMR; Protein DMR; WD repeat-containing protein 20; WD repeat domain 20; anti-WDR20 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	250ug/ml (varies by lot)
Applicable Applications for anti-WDR20 antibody	IHC (Immunohistochemistry)
Predicted Reactivity	Mouse, Rat, Chicken, Turkey, Rabbit, Orangutan, Monkey, Gorilla, Chimpanzee, West Indian ocean coelacanth, Duckbill platypus, Nile tilapia, Crab-eating macaque, White-tufted-ear marmoset, Chinese hamster, Green anole, Zebra finch, Three-spined stickleback, Small-eared galago, Japanese pufferfish, Medaka fish, Thirteen-lined ground squirrel, Northern white-cheeked gibbon
Immunogen	Between 450 and 500
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



IHC (Immunohistochemistry)

Detection of human WDR20 by immunohistochemistry. Sample: FFPE section of human seminoma. Antibody: Affinity purified rabbit anti-WDR20 (Cat. No. AAA214101) used at a dilution of 1:250. Detection: DAB



IHC (Immunohistochemistry)

Detection of human WDR20 by immunohistochemistry. Sample: FFPE section of human thyroid carcinoma. Antibody: Affinity purified rabbit anti-WDR20 (Cat. No. AAA214101) used at a dilution of 1:250. Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-WDR20 antibody	WDR20 (WD repeat-containing protein 20) contains five WD repeats and is a member of the WD family of proteins. The WD repeat is defined by four or more repeating units of a conserved core of approximately 40 amino acids ending with tryptophan-aspartic acid (WD). WD repeats may serve as sites of protein-protein interaction for adaptor proteins and facilitate multiprotein complex formation. WD proteins are involved in a variety of cellular processes. The function of WDR20 has not been characterized.
Product Categories/Family for anti-WDR20 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GeneID	91833
UniProt Accession #	Q8TBZ3
UniProt Protein Name	WD repeat-containing protein 20
UniProt Gene Name	WDR20
UniProt Entry Name	WDR20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/wdr20-antibody-immunohistochemistry-elisa/214101>