

Mouse Thomsen-Friedenreich Antigen/CD176 Monoclonal Antibody

Thomsen-Friedenreich Antigen/CD176 (Pan Carcinoma Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA215331	Mouse	Human, Mouse, Rat	Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

Thomsen-Friedenreich Antigen/CD176; Monoclonal Antibody; Thomsen-Friedenreich Antigen/CD176 (Pan Carcinoma Marker); T-F Antigen; TF Antigen; Asialoglycophorin; pan-carcinoma marker; CD176; anti-Thomsen-Friedenreich Antigen/CD176 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgM, kappa
Clone Number	A78-G/A7
Specificity	Recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF, including related structures on the glycolipid) and shows no cross-reactivity with sialylated glycophorin. The Thomsen-Friedenreich antigen acts as an oncofetal antigen, with low expression in normal adult tissues but increasing to fetal levels of expression in hyperplasia or malignancy. It is considered as a pan-carcinoma marker. This MAb is capable to agglutinate desialylated red blood cells. During metastasis, the ability of malignant cells to form multicellular aggregates via homotypic or heterotypic aggregation and their adhesion to the endothelium are critical. The tumor-associated carbohydrate Thomsen-Friedenreich antigen (Gal-GalNAc) is involved in tumor cell adhesion and tissue invasion. It also causes an immune response, and overexpression of the antigen causes cancer cells to be more sensitive to natural killer cell lysis. The Thomsen-Friedenreich antigen is suppressed in normal healthy cells and represents one of the few chemically well-defined antigens associated with tumor malignancy. The presence of the Thomsen-Friedenreich antigen on the surface of cancer cells may result from a divergence from the normal pathway for O-linked glycosylation in these cells, most likely caused by inappropriate localization of the enzymes involved in synthesis of the disaccharide.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA at 1.0mg/ml
Form/Format	200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Applicable Applications for anti-Thomsen-Friedenreich Antigen/CD176 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence)
Positive Control	KG1 cells. Human colorectal carcinoma tissues.
Immunogen	Neuraminidase-treated human red blood cells
Cellular Localization	Cell Surface
Preparation and Storage	Antibody with azide: store at 2 to 8°C. Antibody without azide: store at -20 to -80°C. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with T-F Antigen/CD176
Mouse Monoclonal Antibody (A78-G/A7).

ADDITIONAL INFORMATION

References	1. Ryder SD et al.Peanut ingestion increases rectal proliferation in individuals with mucosal expression of peanut lectin receptor. Gastroenterology 1998, 114(1):44-9. 2. Karsten U et al. A new monoclonal antibody (A78-G/A7) to the Thomsen-Friedenreich pan-tumor antigen. Hybridoma 1995, 14(1):37-44. 3. Kanitakis J et al. Differential expression of the cancer associated antigens T (Thomsen-Friedenreich) and Tn to the skin in primary and metastatic carcinomas. J Clin Pathol 1998, 51(8):588-92. 4. Baldus SE et al. Coexpression of MUC1 mucin peptide core and the Thomsen-Friedenreich antigen in colorectal neoplasms. Cancer 1998, 82(6):1019-27. 5. Cao Y. et al., Thomsen-Friedenreich-related carbohydrate antigens in normal adult human tissues: a systematic and comparative study. Histochem Cell Biol 106:197-207 (1996).
------------	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/thomsen-friedenreich-antigen-cd176-antibody-clone-a78-g-a7-immunohistochemistry-immunofluorescence-elisa/215331>