

Mouse anti-Human CCD8A Monoclonal Antibody | anti-CD8A antibody

CCD8A (Cytotoxic- & Suppressor T-Cell Marker) Recombinant Mouse Monoclonal Antibody [Clone rCD8/6590]

CATALOG # AAA216044	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

CCD8A; Monoclonal Antibody; CCD8A (Cytotoxic- & Suppressor T-Cell Marker) Recombinant Mouse Monoclonal Antibody [Clone rCD8/6590]; CCD8A (Cytotoxic- & Suppressor T-Cell Marker); CD8 antigen; alpha polypeptide (p32); T8/Leu-2 T-lymphocyte differentiation antigen; Ly3; LYT3; MAL; T-cell surface glycoprotein CD8 alpha chain; anti-CD8A antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG1, kappa
Clone Number	rCD8/6590
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-CD8A antibody	IHC (Immunohistochemistry)
Antibody Type	Recombinant Antibody
Immunogen	Synthetic peptide corresponding to CD8? residues within aa135-235 of CD8?
Cellular Localization	Cell Surface
Positive Control	HuT78 or hPBL cells. Human lymph node or tonsil.
Hu-chromosome Location	2p11.2
Unigene	85258
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human tonsil stained with CD8 Recombinant Mouse Monoclonal Antibody (rCD8/6590).

ADDITIONAL INFORMATION

Related Product Information for anti-CD8A antibody	CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC class 1 and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells. For mature T-cells, CD4 and CD8 are mutually exclusive, so anti-CD8, generally used in conjunction with anti-CD4. It is a useful marker for distinguishing helper/inducer T-lymphocytes, and most peripheral T-cell lymphomas are CD4+/CD8-. Anaplastic large cell lymphoma is usually CD4+ and CD8-, and in T-lymphoblastic lymphoma/leukemia, CD4 and CD8 are often co-expressed. CD8 is also found in littoral cell angioma of the spleen.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	225007536
NCBI GeneID	925
NCBI Accession #	NP_001139345.1
NCBI GenBank Nucleotide #	NM_001145873.1
UniProt Accession #	P01732; Q13970; Q4ZG17; D6W5M8
Molecular Weight	32kDa
NCBI Official Full Name	T-cell surface glycoprotein CD8 alpha chain isoform 1
NCBI Official Synonym Full Names	CD8a molecule
NCBI Official Symbol	CD8A
NCBI Official Synonym Symbols	CD8; MAL; p32; Leu2
NCBI Protein Information	T-cell surface glycoprotein CD8 alpha chain; T8 T-cell antigen; T cell co-receptor; OKT8 T-cell antigen; T-cell antigen Leu2; Leu2 T-lymphocyte antigen; CD8 antigen, alpha polypeptide (p32); T-lymphocyte differentiation antigen T8/Leu-2
UniProt Protein Name	T-cell surface glycoprotein CD8 alpha chain
UniProt Gene Name	CD8A
UniProt Synonym Gene Names	MAL
UniProt Entry Name	CD8A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/ccd8a-antibody-clone-rcd8-6590-immunohistochemistry-elisa/216044>