

Human Interferon Alpha/Beta Receptor 2 (IFNa/bR2) RTU ELISA Kit | IFNa/bR2 rtu elisa kit

Human Interferon Alpha/Beta Receptor 2 (IFNa/bR2) ELISA Kit

CATALOG #
AAA216145

REACTIVITY
Human


PRODUCT SYNONYMS

Interferon Alpha/Beta Receptor 2 (IFNa/bR2); Human Interferon Alpha/Beta Receptor 2 (IFNa/bR2) ELISA Kit; IFNa/bR2; IFN-alpha-REC; IFN-R; IFNABR; IFNARB; Interferon(Alpha; Beta And Omega)Receptor Beta Chain; Interferon alpha binding protein; Type I interferon receptor 2; IFNa/bR2 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Assay Type	Sandwich
Samples	Serum, Plasma, Urine or Other Biological Fluids.
Detection Range	0.625-40ng/mL
Sensitivity	0.235ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IFNa/bR2 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IFNa/bR2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IFNa/bR2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IFNa/bR2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of IFNa/bR2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.