

Mouse Stromal Cell Derived Factor 1 (SDF1) RTU ELISA Kit | SDF1 rtu elisa kit

Mouse Stromal Cell Derived Factor 1 (SDF1) ELISA Kit

CATALOG #
AAA218328

REACTIVITY
Mouse


PRODUCT SYNONYMS

Stromal Cell Derived Factor 1 (SDF1); Mouse Stromal Cell Derived Factor 1 (SDF1) ELISA Kit; CXCL12; PBSF; P-BSF; SCYB12; SDF1-A; SDF1-B; SDF1A; SDF1B; TLSF-A; TLSF-B; TPAR1; Pre-B Cell Growth-Stimulating Factor; Chemokine C-X-C-Motif Ligand 12; SDF1 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	93
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.059ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SDF1 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SDF1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SDF1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SDF1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of SDF1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	7305465
NCBI GeneID	20315
NCBI Accession #	NP_038683.1
NCBI GenBank Nucleotide #	NM_013655.4
UniProt Accession #	P40224; Q4FJL5; Q543V6
Molecular Weight	10,032 Da
NCBI Official Full Name	stromal cell-derived factor 1 isoform beta
NCBI Official Synonym Full Names	chemokine (C-X-C motif) ligand 12
NCBI Official Symbol	Cxcl12
NCBI Official Synonym Symbols	Pbsf; Sdf1; Tlsf; Tpar1; Scyb12
NCBI Protein Information	stromal cell-derived factor 1
UniProt Protein Name	Stromal cell-derived factor 1
UniProt Gene Name	Cxcl12
UniProt Synonym Gene Names	Sdf1; SDF-1; TPAR1; PBSF; TLSF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-rtu-elisa-kits/stromal-cell-derived-factor-1-sdf1/218328>