

Rabbit anti-Human MMP9 Polyclonal Antibody | anti-MMP9 antibody

Rabbit anti-MMP9 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220331	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MMP9; Polyclonal Antibody; Rabbit anti-MMP9 Antibody; CLG4B; Matrix metalloproteinase-9; MMP-9; 92 kDa gelatinase; 92 kDa type IV collagenase; Gelatinase B; GELB; anti-MMP9 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-MMP9 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MMP9. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MMP9 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MMP9 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MMP9 expression in HT1080 (A), HuvEc (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MMP9 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	74272287
NCBI GeneID	4318
NCBI Accession #	NP_004985.2
NCBI GenBank Nucleotide #	NM_004994.2
UniProt Accession #	P14780; Q3LR70; Q8N725; Q9H4Z1; Q9UCJ9; Q9UCL1; Q9UDK2; B2R7V9
Molecular Weight	78,458 Da
NCBI Official Full Name	matrix metalloproteinase-9 preproprotein
NCBI Official Synonym Full Names	matrix metallopeptidase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)
NCBI Official Symbol	MMP9
NCBI Official Synonym Symbols	GELB; CLG4B; MMP-9; MANDP2
NCBI Protein Information	matrix metalloproteinase-9; 92 kDa gelatinase; type V collagenase; macrophage gelatinase; 92 kDa type IV collagenase; matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)
UniProt Protein Name	Matrix metalloproteinase-9
UniProt Gene Name	MMP9
UniProt Synonym Gene Names	CLG4B; MMP-9; GELB
UniProt Entry Name	MMP9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/mmp9-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220331>